

Assessment of vulnerable plaque by OCT

Comparison among Intravascular Imaging Modalities

Takashi Akasaka, M.D.

Department of Cardiovascular Medicine

Wakayama Medical University

Wakayama, Japan





Disclosure Statement of Financial Interest

I (Takashi Akasaka) do not have a financial interest/arrangement or affiliation with one or more organizations that could be perceived as a real or apparent conflict of interest in the context of the subject of this presentation.



Pathohistology of vulnerable plaque (HE stain)



Thin fibrous cap
Large lipid core
Advanced atherosclerosis




Pathohistological characteristics of vulnerable plaque

- Positive remodeling
- Eccentric plaques
- Lipid-rich plaques (necrotic core)
- Thin fibrous cap ($< 65 \mu\text{m}$)
- Rupture (60%) or ulceration (30~40%) of fibrous caps
- Thrombus formation
- Macrophage accumulation



Identification of vulnerable plaque

- Plaque prone to rupture



Study Population

Inclusion criteria

Acute Myocardial Infarction

continuous chest pain lasted > 30 minutes,

arrival within 6 hours from the onset of chest pain

ST elevation ≥ 0.1 mV in 2 or more contiguous leads

culprit lesion with diameter stenosis $\geq 75\%$

TIMI flow ≤ 2 identified by CAG

Exclusion criteria

left bundle-branch block

pacemaker rhythm

a culprit lesion in the left main coronary artery

history of prior MI

cardiogenic shock

unsuccessful reperfusion < TIMI III flow by thrombectomy



Study Design

Oral aspirin (162 mg) and intravenous heparin (100 U/kg) were administered before PCI.

Cardiac catheterization was performed by the femoral approach, using a 7F sheath and catheters.

- **Thrombectomy** (*Export catheter* © Medtronic Japan)

TIMI grade III


- **IVUS** (*Atlantis SR Pro* © 2.5F, 40-MHz; Boston Scientific, Natick, MA, USA)

- **CAS** (*Angioscope MC-800E and the optic fiber AS-003*, Nihon Kohden)

- **OCT** (*ImageWire* ©; LightLab Imaging, Westford, MA, USA)



OCT system (LightLab Co.)



PIU

Wakayama Medical University




Optical Coherence Tomography (OCT)





OCT vs histology

Fibrous




**Homogeneous,
Signal-rich**

Lipid pool



**Echolucent,
Diffuse Borders**

Calcific



**Echolucent,
Sharp Borders**



Tissue characterization by OCT and IVUS

Kume T, Akasaka T, et al (Am J Cardiol 97 : 1172 - 1175, 2006)



Pathohistological Diagnosis	Sensitivity	Specificity	Predictive Value	
			Positive	Negative
OCT image				
Fibrous (n=43)	79	99	97	93
Fibrocalcific (n=82)	96	88	89	96
Lipid (n=41)	85*	94	83	95
IVUS image				
Fibrous (n=43)	88	86	69	95
Fibrocalcific (n=82)	98	96	96	98
Lipid (n=41)	59	97	86	87

Data are demonstrated as percentages. * $p < 0.05$ vs IVUS

Wakayama Medical University






Comparison between IVUS and OCT

	IVUS	OCT
<i>Scan area</i>		
<i>Max. penetration depth</i>	4 - 8 mm	1 - 1.5 mm
<i>Blood clearing To obtain images</i>	Not required	Required



Pathohistological characteristics of vulnerable plaque




Assessment by OCT

- Positive remodeling 
- Eccentric plaques 
- Lipid-rich plaques (necrotic core) 
- Thin fibrous cap
- Rupture or ulceration of fibrous caps
- Thrombus formation
- Macrophage accumulation




Pathohistological characteristics of vulnerable plaque

Assessment by OCT


- Positive remodeling 
- Eccentric plaques 
- Lipid-rich plaques (necrotic core) 
- Thin fibrous cap
- Rupture or ulceration of fibrous caps
- Thrombus formation
- Macrophage accumulation



Inferior AMI (71y.o. Male)




Inferior-AMI (71y.o., M) Plaque Rupture



→ Ruptured Fibrous Cap
↕ Fibrous Cap Thickness = 40 μm


TL : True Lumen

UL : Ulceration



Ruptured Fibrous Caps


OCT




CAS



IVUS



Anteroseptal AMI (80y.o., M)



Fibrous cap thickness = 60 μm



Wakayama Medical University






Thickness of fibrous caps

Histology vs OCT



Kume T, Akasaka T, et al (Am Heart J . 152:755, 2006)



Anteroseptal AMI (80 y.o. Male)



Anteroseptal AMI (80y.o., M)



↑ Erosion (Ulceration)


↑ Thrombus





Inf-AMI (71y.o., M)

Thrombus

Red Thrombus




Thrombectomy



Intensity Half Distance
= 135 μm



Thrombus



OCT

CAS

IVUS





Differentiation between red and white thrombus

Peak intensity



Intensity half distance

Red thrombus



183 ± 42

White thrombus



324 ± 50 *

* $p = 0.0001$

Kume T, Akasaka T, et al (Am J Cardiol . 2006, 97:1713-1717)

Wakayama Medical University





Differentiation between red and white thrombus

White thrombus



Red thrombus



Comparison of plaque Images in AMI (OCT vs. CAS vs. IVUS) n=30

(Kubo T, Akasaka T, et al. J Am Coll Cardiol 50:933-999,2007)

	OCT	*CAS	**IVUS	*p	**p
Plaque Rupture (%)	73	47	40	0.035	0.009
Ulceration (erosion) (%)	23	3	0	0.022	0.005
Thrombus (%)	100	100	33	1.000	<0.001
Red thrombus (%)	100	90	-	0.076	-
White thrombus (%)	100	93	-	0.150	-
TCFA ($\leq 65 \mu m$) (%)	83	-	-	-	-
Fibrous cap thickness (μm)	49 \pm 21	-	-	-	-
LRP (Lipid Arch $> 180^\circ$) (%)	83	-	67	-	NS









TCFA; Thin Cap Fibro-Atheroma, LRP; Lipid Rich Plaque

Wakayama Medical University

Pathohistological characteristics of vulnerable plaque

Assessment by OCT

- Positive remodeling 
- Eccentric plaques 
- Lipid-rich plaques (necrotic core) 
- Thin fibrous cap 
- Rupture or ulceration of fibrous caps 
- Thrombus formation 
- Macrophage accumulation





OCT findings


Low M ϕ

High M ϕ


OCT




CD68
(macrophage)



Macrophages










Macrophages



Pathohistological characteristics of vulnerable plaque

Assessment by OCT








- Positive remodeling 
- Eccentric plaques 
- Lipid-rich plaques (necrotic core) 
- Thin fibrous cap 
- Rupture or ulceration of fibrous caps 
- Thrombus formation 
- Macrophage accumulation 



Summary


Assessment of vulnerable plaque by OCT

Comparison with pathohistology

- Positive remodeling 
- Eccentric plaques 
- Lipid-rich plaques (necrotic core) 
- Thin fibrous cap 
- Rupture or ulceration of fibrous caps 
- Thrombus formation 
- Macrophage accumulation 



Corresponding Images of OCT and Angioscopy




(Kubo T, et al. J Am Coll Cardiol Interv 1:74-80,2008)

Figure 1
Wakayama Medical University




Angioscopy vs OCT

Plaque color vs lipid size



Plaque color vs fibrous cap thickness




(Kubo T, et al. J Am Coll Cardiol Intv 1:74-80,2008)

Wakayama Medical University



IVUS-derived TCFA

(Rodriguez-Granillo GA, et al. J Am Coll Cardiol 46:2038-2042, 2005)




- ◆ Percent atheroma volume = $(\text{EEM area} - \text{Lumen area}) / \text{EEM area} \times 100 \geq 40\%$
- ◆ Necrotic core $\geq 10\%$
- ◆ Without evident overlying fibrous tissue




VH-IVUS vs OCT


Figure 2



Without evident overlying fibrous tissue




Without evident overlying fibrous tissue



With evident overlying fibrous tissue



Distribution of fibrous cap thickness in cases with VH-TCFA



Concordance & discordance between VH-IVUS and OCT in the assessment of TCFA

Table 4

IVUS-VH Diagnosis \ OCT Diagnosis	TCFA (n=11)	Not TCFA (n=36)
VH-TCFA (n=31)	9	22
Not VH-TCFA (n=16)	2	14



Wakayama Medical University




Summary

- **OCT can identify lipid-rich plaques more sensitively compared with IVUS.**
- **OCT can demonstrate rupture or ulceration of fibrous cap with higher detection rate than that of IVUS or CAS.**
- **OCT could detect intracoronary thrombus almost exclusively which was confirmed by CAS.**
- **OCT may estimate macrophage accumulation within fibrous caps.**



Frequency domain OCT (M4)




Conclusion


OCT should be the most ideal tool in the assessment of vulnerable plaques among various intravascular imaging modalities .



Unstable AP




Three vessel OCT examinations in a patient with posterior AMI



The culprit lesion was LCX (#11), and TCFA (①), plaque rupture (②,③) and intracoronary thrombus (②, ③, ④) were observed by OCT. Although the plaques in LAD (⑦, ⑧) were not observed, plaque rupture (⑤, ⑥) were detected by OCT in the non-culprit lesions of LCX (#13).



Three vessel OCT examinations in a patient with posterior AMI



Also OCT revealed TCFA (⑨-⑬) and plaque rupture(⑪,⑫,⑬) in the non-culprit lesions of RCA.



OCT analysis of the non-culprit plaques

	Non-culprit plaque		p-value
	AMI (n=43)	SAP (n=25)	
Plaque rupture (n, %)	10 (24)	1 (4)	0.035
Plaque ulceration (n, %)	1 (2)	0 (0)	0.632
Intracoronary thrombus (n, %)	11(26)	0 (0)	0.006
Fibrous cap thickness (μ m)	109 \pm 55.5	194 \pm 81.9	<0.001
Lipid arc > 90° (n, %)	18 (42)	12 (48)	0.623
TCFA (n, %)	15(35)	2 (8)	0.012




Pathohistological characteristics of vulnerable plaque

- Positive remodeling
- Eccentric plaques
- Lipid-rich plaques (necrotic core)
- Thin fibrous cap
- Rupture or ulceration of fibrous caps
- Thrombus formation
- Macrophage accumulation



Thrombus




OCT

CAS

IVUS






Optical Coherence Tomography (OCT)





- Size of imaging core (0.4 mm)
- Microscopic resolution (10-20 μ m)
- Real time Imaging (15 frames/s)





Fibrous plaque




Fibrocalcific plaque




Fibro-lipidic plaque



Comparison between IVUS and OCT



IVUS



OCT

Resolution	(axial) 100 - 150 μm (lateral) 150 - 300 μm	15 - 20 μm 25 - 40 μm
Size of imaging core	0.8 mm	0.4 mm
Dynamic range	40 - 60 dB	90 - 110 dB
Frame rate	30 frames/s	15 frames/s

