Renal Artery Stenosis and Abdominal Aortic Aneurysm

DR. Rabin Chakraborty


MD, DNB, FRCP(London), FRCP(Glasgow), FICC, FISE, FCSI, DM (Cardiology)

Director and Head, Senior Consultant Interventional cardiologist Apollo *Gleneagles* Heart Institutes, Kolkata, INDIA

- DKS 62 yrs / Male
- Hypertension, Euglycemic, Heavy smoker
- C/C: Abdominal pain (dull aching) for last 2-3 months.
 - Prominent pulsation in lower abdomen for same duration.
- Clinical examination revealed large pulsatile mass in left lower abdomen.
- No Hx of Angina


Contrast enhanced CT scan shows large (10.5×6.5 cm) infrarenal aortic aneurysm





Contrast enhanced CT scan shows large (10.5×6.5 cm) infrarenal aortic aneurysm






Abdominal aortogram shows large infrarenal aneurysm with right renal artery stenosis




PLAN

- Right renal angioplasty
- Main body of the abdominal aortic bifurcated stent graft from left femoral artery
- Contra-lateral limb from right femoral artery
- We selected AAA stent graft from Medtronic (Endurant stent graft)


Endurant bifurcated stent graft




Endurant stent graft



Left radial approach, 6F JR3.5, BMW wire, balloon 4.0 × 15 mm



Right Renal Angioplasty from Left Radial Artery



Balloon Dilatation



Hippocampus 7.0×15 mm




Hippocampus 7.0×15 mm




Hippocampus 7.0×15 mm

DIPAK KUMAR SARKAR 52Y/M DR.RABIN CHAKRABORTY*** APOLLO GLENEAGLES HOSPITALS, KOLKATA 27800/189/2012 20120112 180027.549 Lao - 11 deg "Cran - 0 deg Run 10 Of 43 Zoom: 99% Frame 1 Of 29

Main body: Endurant 32×160 mm






Not yet deployed fully




Proximal Fixation



6F Pigtail through the stent across the opening for contra-lateral limb from left radial



Snaring of the retrograde wire







Contra-lateral limb Endurant 10×124 mm




Contra-lateral limb Endurant 10×124 mm



Final deployment of the main body




Post Deployment Injection: Diseased left common iliac artery distal to the stent




Low pressure dilatation of ipsilateral limb 10×40 mm balloon



Extension of ipsilateral limb with Endurant 10×82 mm



Low pressure dilatation of whole stent including overlapping areas with a compliant Reliant stent graft balloon.



Low pressure dilatation of whole stent including overlapping areas with a compliant Reliant stent graft balloon



Low pressure dilatation of whole stent including overlapping areas with a compliant Reliant stent graft balloon.




Final result



Procedure summary

- Procedure date 12.01.2012.
- Under GA (Hybrid with CTU Surgeon)
- Surgical exposure & closure of both femoral arteries
- Total procedure time 95 mins.
- Total UFH 9,000 IU.
- Total contrast used 400 ml Omnipaque.
- Extubated after 1 hr post procedure.
- No blood transfusion required.
- Planning follow up imaging by contrast CT after 6 m



Thank You