

Update and Insight from Plaque Imaging PREVENT Trial

Seung-Jung Park, MD, PhD

Professor of Medicine, University of Ulsan College of Medicine,
Heart Institute, Asan Medical Center, Seoul, Korea

Vulnerable Plaque

To Treat or Not To Treat ?



Thin-cap Fibroatheroma (TCFA)

Rupture/Healed Rupture

Erosion

Erosion/Thrombus

Calcific Nodule



Thin-cap Fibroatheroma (TCFA)


Rupture/Healed Rupture

Confluent Necrotic Core

>50% Area Narrowing

Calcium >5%





55 y/o male, Effort Chest Pain



Plaque Characteristics by OCT, VH-IVUS & NIRS


Rupture, TCFA

$\max LCBI_{4mm} = 404$



Necrotic Core 25%

Functionally Insignificant To Treat or Not To Treat? Vulnerable Plaque



FFR : 0.89

Angiographic DS : **70%**

IVUS MLA : **3.45 mm²**

Plaque burden : **73%**

max LCBI_{4mm} : **404**

TCFA (+)

Not to Treat ?

Negative FFR (non-invasive stress tests) means *just excellent prognosis (0.6%/year, Cardiac Death and MI)*, even in the presence of angiographically proven coronary artery disease.


Shaw LJ, J Nucl Cardiol 2004;11:171-85 ,Prognostic value of gated myocardial perfusion SPECT. Very large meta-analysis (n=39,173 patients)

To Treat ?

Vulnerable Plaque (defined by PROSPECT study) *has more tendency to increase MACE.*

PROSPECT: MACE

(N=700, ACS, 3-Vessel Imaging after PCI)




Number at risk

	0	1	2	3
ALL	697	557	506	480
CL related	697	590	543	518
NCL related	697	595	553	521
Indeterminate	697	634	604	583

Vulnerable Plaque Defined by VH-IVUS

Independent Predictors of Non-Culprit Lesion Events



$PB_{MLA} \geq 70\%$


VH-TCFA

$MLA \leq 4.0 \text{ mm}^2$

	HR [95% CI]	P value
$PB_{MLA} \geq 70\%$	5.03 [2.51, 10.11]	<0.0001
VH-TCFA	3.35 [1.77, 6.36]	0.0002
$MLA \leq 4.0 \text{ mm}^2$	3.21 [1.61, 6.42]	0.001

Stone GW et al. NEJM 2011;364:226-35

PROSPECT: Correlates of Non Culprit Lesion Related Events



*Likelihood of one or more such lesions being present per patient. PB = plaque burden at the MLA

Q1,

**Can *Optimal Medical Treatment*
Stabilize *Plaque Vulnerability* ?**

STABLE Trial

(STatin and Atheroma VulneraBiLity Evaluation)

Double-blinded, Prospective, Randomized, Controlled Trial

290 patients with
Deferred native coronary artery lesion

2:1 randomization, double-blinded


Rosuvastatin 40mg


Rosuvastatin 10mg

Primary efficacy endpoint; Change in %NC volume
within target segment by VH-IVUS at 1 year

Secondary endpoint: change in %NC volume comparing rosuvastatin
40mg vs. 10mg.

Primary Endpoint %NC Volume Changes at 1 Year





	Baseline	1 year
EEM, mm ²	19.0	14.0
Plaque, mm ²	14.6	10.3
Lumen, mm ²	4.4	3.7
VH-%NC	30%	15%
VH-TCFA	+	-
OCT-TCFA	+	-

Rosuvastatin Treatment Can Make A Plaque Regression and Stabilization

Clinical Outcomes at 1 Year

- No cardiac death
- Culprit-related MACE: 4 pts (2.3%).
- Non Culprit-related MACEs: 8 pts (3.6%).
- No Difference in Non Culprit-MACE between rosuvastatin 40 vs.10mg (3.9 vs. 2.7%, p=NS)

Can **Optimal Medical Treatment** Stabilize Plaque Vulnerability ?


***Yes, Rosuvastatin Therapy Can Make A
Plaque Regression and Stabilization.***

Q2,

Can *BVS*

***Stabilize Plaque Vulnerability and
Make an Any Difference ?***

*BVS Can Make **Plaque** Stabilization and Lumen Enlargement*



Pre-PCI

Post-PCI


6 months

2 years


5 years

Vessel area (mm ²)	15.72	15.34 (3%)	14.09 (10%)	13.76 (12%)
Plaque area (mm ²)	8.78	9.17 (4%)	7.54 (14%)	7.07 (19%)
Mean LA (mm ²)	6.95	6.17 (11%)	6.56 (5.6%)	8.09 (16%)

ABSORB II, 1-year Results




ABSORB III, 1-year Results




What's the Difference ?

Optimal Medical
Treatment



Stabilized Plaque Vulnerability
Decreased Plaque Volume
Decrease Vessel Size
Decreased Lumen


BVS




Stabilized Plaque Vulnerability
Decreased Plaque Volume
Decrease Vessel Size
Increased Lumen

Pre-procedural IVUS-VH and Vessel Remodeling over 3 years in ABSORB II

Absorb (n=224)




Xience (n=123)




Various balloon-artery ratios and vessel remodeling

Absorb (n=224)



Xience (n=123)



PREVENT Study,

The **PREVENT**ive Implantation of BVS on Stenosis With Functionally Insignificant Vulnerable Plaque Compared to Optimal Medical treatment.

Objective,





To determine whether BVS implantation on functionally insignificant vulnerable plaque, reduce the incidence of the composite of MACEs compared with optimal medical therapy alone.

A prospective, randomized, multicenter, clinical trial with 'all comers' design. Approximately 2,000 patients will be enrolled from international heart centers.

Defining, Functionally Insignificant Vulnerable Plaque



1. $PB_{MLA} \geq 70\%$
2. $MLA \leq 4.0 \text{ mm}^2$
3. TCFA by OCT or VH-IVUS
4. LRP on NIRS ($_{\max}LCBI_{4\text{mm}} > 315$)



PREVENT Trial

Any Epicardial Coronary Stenosis (≤ 40 mm) with FFR ≥ 0.80 and with Two of the following

1. Plaque Burden $>70\%$
2. MLA $\leq 4.0\text{mm}^2$
3. TCFA by OCT or VH-IVUS
4. Lipid-Rich Plaque on NIRS ($_{\text{max}}\text{LCBI}_{4\text{mm}} > 315$)

R

BVS+OMT
N=800

OMT
N=800

Primary endpoint *at 2 years*:
CV death, MI, Hospitalization d/t unstable angina

OCT sub-study/ NIRS sub-study, (300 patients in each arm at 2 years)

Inclusion Criteria

Age 18 years or older,
Symptomatic or asymptomatic coronary stenosis,
Eligible lesions for PCI (≤ 40 mm), with
FFR >0.80 and met the two of the following

1. Plaque burden $>70\%$
2. MLA $<4\text{mm}^2$
3. TCFA by OCT or VH-IVUS
4. Lipid-rich plaque on NIRS ($_{\max}\text{LCBI}_{4\text{mm}} > 315$)


Primary and Major Secondary End Point,

The primary endpoint is the 2-year MACE (cardiovascular death, nonfatal MI, unplanned rehospitalization due to unstable angina).

The secondary endpoints include overall MACE, non-urgent revascularization, and rate of cerebrovascular event.


BVS cases

M/58, Unstable Angina






FFR

Intravenous adenosine, 140 $\mu\text{g}/\text{kg}/\text{min}$






Imaging

MLA : 2.7 mm²
Plaque burden 77 %




Thrombi & Erosion



Randomized to BVS

58 y/o male, Unstable Angina



Angiographic DS : 50%

FFR : 0.81



IVUS MLA : 2.7 mm²

Plaque burden : 77 %


Erosion (+)

max LCBI_{4mm} : 0


BVS




Pre-Dilate, NC
3.0 mm x 15 mm



Absorb BVS
3.5 mm x 18 mm




After BVS




NC Balloon,
4.0 mm x 13 mm

BVS



OCT Confirmed Good Apposition of BVS

55 y/o male,
Unstable Angina



Culprit PCI for RCA and LM-pLAD

Pre



Xience Alpine
3.0 x 18 mm


Xience Alpine
4.0 x 23 mm

Post
PCI




Non-Culprit LCX, Randomized to OMT Group


MLA 2.8 mm²
Plaque burden 81%



maxLCBI4mm : 930




**11 months later,
Recurred Chest Pain**




Disease Progression !

OMT group, PCI






Xience Alpine
3.5 x 23mm

57 y/o Female,
Atypical Chest Pain




Vulnerable Plaque by OCT & NIRS



MLA 2.7 mm²
Plaque burden 73%

maxLCBI 4mm : 571

Randomized to OMT



FFR : 0.85

Angiographic DS : 50%

IVUS MLA : 2.7 mm²



Plaque burden : 73%

max LCBI_{4mm} : 571

TCFA (+)


Functionally
Insignificant
Vulnerable Plaque

7 months later,
Rest Chest Pain



Disease Progression !

OMT group, PCI




Resolute Onyx

3.5 x 18 mm

2.5 x 15 mm

Current Patients Enrollment 2017 Mar.





Thank You !!

summitMD.com