

"Full metal jacket" (stented length  $\geq$  64 mm) using drug-eluting stents for de novo coronary artery lesions J. Aoki, A. T. Ong, G. A. Rodriguez Granillo, E. P. McFadden, C. A. van Mieghem, M. Valgimigli, K. Tsuchida, G. Sianos, E. Regar, P. P. de Jaegere, W. J. van der Giessen, P. J. de Feyter, R. T. van Domburg and P. W. Serruys  
Am Heart J (2005) 150;994-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16290984](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16290984)

Comparison of sirolimus-eluting stent, paclitaxel-eluting stent, and bare metal stent in the treatment of long coronary lesions Y. H. Kim, S. W. Park, C. W. Lee, M. K. Hong, H. C. Gwon, Y. Jang, M. M. Lee, B. K. Koo, D. J. Oh, K. B. Seung, S. J. Tahk, J. Yoon and S. J. Park

Catheter Cardiovasc Interv (2006) 67;181-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16400663](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16400663)