

Sirolimus-eluting stents versus bare-metal stents in patients with in-stent restenosis: results of a pooled analysis of two randomized studies F. Alfonso, M. J. Perez-Vizcayno, R. Hernandez, C. Fernandez, J. Escaned, C. Banuelos, A. Bethencourt, J. R. Lopez-Minguez, J. Angel, A. Cequier, M. Sabate, C. Moris, J. Zueco and R. Seabra-Gomes *Catheter Cardiovasc Interv* (2008) 72; 459-67
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18814274

Early stent thrombosis in patients with acute coronary syndromes treated with drug-eluting and bare metal stents: the Acute Catheterization and Urgent Intervention Triage Strategy trial J. Aoki, A. J. Lansky, R. Mehran, J. Moses, M. E. Bertrand, B. T. McLaurin, D. A. Cox, A. M. Lincoff, E. M. Ohman, H. D. White, H. Parise, M. B. Leon and G. W. Stone *Circulation* (2009) 119; 687-98
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19171852

Late outcomes of drug-eluting versus bare metal stents in saphenous vein grafts: Propensity score analysis R. J. Applegate, M. Sacrinty, M. Kutcher, R. Santos, S. Gandhi and W. Little *Catheter Cardiovasc Interv* (2008) 72; 7-12
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18412239

"Off-label" stent therapy 2-year comparison of drug-eluting versus bare-metal stents R. J. Applegate, M. T. Sacrinty, M. A. Kutcher, R. M. Santos, S. K. Gandhi, T. T. Baki and W. C. Little *J Am Coll Cardiol* (2008) 51; 607-14
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18261678

Incidence of coronary stent thrombosis based on academic research consortium definitions R. J. Applegate, M. T. Sacrinty, W. C. Little, R. M. Santos, S. K. Gandhi and M. A. Kutcher *Am J Cardiol* (2008) 102; 683-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18773988

Comparison of outcomes of drug-eluting stents versus bare-metal stents in nonstial proximal left anterior descending coronary arteries L. Bonello, A. De Labriolle, G. Lemesle, P. Roy, D. H. Steinberg, T. L. Slottow, Z. Xue, R. Torguson, K. Kaneshige, W. O. Suddath, L. F. Satler, K. M. Kent, J. Lindsay, A. D. Pichard and R. Waksman *Am J Cardiol* (2009) 103; 496-500
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195509

A randomized controlled trial of a paclitaxel-eluting stent versus a similar bare-metal stent in saphenous vein graft lesions the SOS (Stenting of Saphenous Vein Grafts) trial E. S. Brilakis, C. Lichtenwalter, J. A. de Lemos, M. Roesle, O. Obel, D. Haagen, B. Saeed, C. Gadiparthi, J. K. Bissett, R. Sachdeva, V. V. Voudris, P. Karyofillis, B. Kar, J. Rossen, P. Fasseas, P. Berger and S. Banerjee *J Am Coll Cardiol* (2009) 53; 919-28
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19281920

Outcomes with drug-eluting stents versus bare metal stents in acute ST-elevation myocardial infarction: results from the Strategic Transcatheter Evaluation of New Therapies (STENT) Group B. R. Brodie, T. Stuckey, W. Downey, A. Humphrey, M.

Nussbaum, S. Laurent, B. Bradshaw, C. Metzger, J. Hermiller, F. Krainin, S. Juk, B. Cheek, P. Duffy and C. A. Simonton Catheter Cardiovasc Interv (2008) 72; 893-900
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19016465

Angiographic and clinical outcome of invasively managed patients with thrombosed coronary bare metal or drug-eluting stents: the OPTIMIST study F. Burzotta, A. Parma, C. Pristipino, A. Manzoli, F. Belloni, G. Sardella, S. Rigattieri, A. Danesi, P. Mazzarotto, F. Summaria, E. Romagnoli, F. Prati, C. Trani and F. Crea Eur Heart J (2008) 29; 3011-21

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18987096

Long-term clinical benefit of drug-eluting stents over bare-metal stents in diabetic patients with de novo left main coronary artery disease: results from a real-world multicenter registry D. Capodanno, M. E. Di Salvo, T. Palmerini, I. Sheiban, M. Margheri, G. Vecchi, G. Sangiorgi, G. Piovaccari, A. Bartorelli, C. Briguori, D. Ardissino, F. Di Pede, A. Ramondo, L. Inglese, A. S. Petronio, L. Bolognese, A. Benassi, C. Palmieri, V. Filippone, S. De Servi and C. Tamburino Catheter Cardiovasc Interv (2009) 73; 310-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19214964

A randomized comparison of sirolimus-eluting versus bare metal stents in the treatment of diabetic patients with native coronary artery lesions: the DECODE study C. Chan, R. Zambahari, U. Kaul, C. P. Lau, H. Whitworth, S. Cohen and M. Buchbinder Catheter Cardiovasc Interv (2008) 72; 591-600

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18949772

Angiographic and clinical outcomes of drug-eluting versus bare metal stent deployment in the Occluded Artery Trial V. Dzavik, C. E. Buller, G. Devlin, R. G. Carere, G. B. Mancini, W. J. Cantor, P. E. Buszman, J. M. Rankin, C. Vozzi, J. R. Ross, S. Forman, B. A. Barton, A. G. Lamas and J. S. Hochman Catheter Cardiovasc Interv (2008)

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19309733

Trends in acute reperfusion therapy for ST-segment elevation myocardial infarction from 1999 to 2006: we are getting better but we have got a long way to go K. A. Eagle, B. K. Nallamothu, R. H. Mehta, C. B. Granger, P. G. Steg, F. Van de Werf, J. Lopez-Sendon, S. G. Goodman, A. Quill and K. A. Fox Eur Heart J (2008) 29; 609-17

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18310671

Balancing the risks of restenosis and stent thrombosis in bare-metal versus drug-eluting stents: results of a decision analytic model P. Garg, D. J. Cohen, T. Gaziano and L. Mauri J Am Coll Cardiol (2008) 51; 1844-53

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18466798

Drug-eluting or bare-metal stenting in patients with diabetes mellitus: results from the Massachusetts Data Analysis Center Registry P. Garg, S. L. Normand, T. S. Silbaugh, R. E. Wolf, K. Zelevinsky, A. Lovett, M. R. Varma, Z. Zhou and L. Mauri Circulation (2008) 118; 2277-85, 7p following 85

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19001019

Lack of clinical long-term benefit with the use of a drug eluting stent compared to use of a bare metal stent in saphenous vein grafts G. Gioia, A. Benassi, R. Mohendra, K. Chowdhury, I. Masood and W. Matthai *Catheter Cardiovasc Interv* (2008) 72; 13-20

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18561143

10-year follow-up of a prospective randomized trial comparing bare-metal stenting with internal mammary artery grafting for proximal, isolated de novo left anterior coronary artery stenosis the SIMA (Stenting versus Internal Mammary Artery grafting) trial J. J. Goy, U. Kaufmann, M. Hurni, S. Cook, F. Versaci, P. Ruchat, O. Bertel, M. Pieper, B. Meier, L. Chiarello and E. Eeckhout *J Am Coll Cardiol* (2008) 52; 815-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18755343

Drug-eluting compared with bare-metal coronary stents among elderly patients P. W. Groeneveld, M. A. Matta, A. P. Greenhut and F. Yang *J Am Coll Cardiol* (2008) 51; 2017-24

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18498954

The impact of lesion length and vessel size on outcomes after sirolimus-eluting stent implantation for in-stent restenosis S. Habara, K. Mitsudo, T. Goto, K. Kadota, S. Fujii, H. Yamamoto, H. Kato, S. Takenaka, Y. Fuku, S. Hosogi, A. Hirono, K. Yamamoto, H. Tanaka, D. Hasegawa, Y. Nakamura, H. Tasaka, S. Otsuru, Y. Okamoto, C. Yamada, M. Miyamoto and K. Inoue *Heart* (2008) 94; 1162-5

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18032460

Comparison of long-term efficacy of the paclitaxel-eluting stent versus the bare-metal stent for treatment of unprotected left main coronary artery disease Y. Han, S. Wang, Q. Jing, Y. Li, H. Liu, Y. Ma, Z. Wang, D. Wang, B. Luan, G. Wang and T. Chen *Am J Cardiol* (2009) 103; 194-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19121435

Late stent malapposition risk is higher after drug-eluting stent compared with bare-metal stent implantation and associates with late stent thrombosis A. K. Hassan, S. C. Bergheanu, T. Stijnen, B. L. van der Hoeven, J. D. Snoep, J. W. Plevier, M. J. Schalij and J. W. Jukema *Eur Heart J* (2009)

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19158118

Neointimal regression-induced incomplete coverage of a bare-metal stent in the left main trunk: Serial angiographic and angioscopic evidence obtained by 5-year follow-up M. Ichikawa and M. Mishima *Catheter Cardiovasc Interv* (2008)

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19309734

Sleep quality and elevated blood pressure in adolescents S. Javaheri, A. Storfer-Isser, C. L. Rosen and S. Redline *Circulation* (2008) 118; 1034-40

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list_uids=18711015

2-year clinical outcomes after implantation of sirolimus-eluting, paclitaxel-eluting, and bare-metal coronary stents: results from the WDHR (Western Denmark Heart Registry)

A. Kaltoft, L. O. Jensen, M. Maeng, H. H. Tilsted, P. Thayssen, M. Bottcher, J. F. Lassen, L. R. Krusell, K. Rasmussen, K. N. Hansen, L. Pedersen, S. P. Johnsen, H. T. Sorensen and L. Thuesen *J Am Coll Cardiol* (2009) 53; 658-64

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19232897

Drug-eluting versus bare metal stents in patients with st-segment-elevation myocardial infarction: eight-month follow-up in the Drug Elution and Distal Protection in Acute Myocardial Infarction (DEDICATION) trial H. Kelbaek, L. Thuesen, S. Helqvist, P. Clemmensen, L. Klovgaard, A. Kaltoft, B. Andersen, H. Thuesen, T. Engstrom, H. E. Botker, K. Saunamaki, L. R. Krusell, E. Jorgensen, H. H. Hansen, E. H. Christiansen, J. Ravkilde, L. Kober, K. F. Kofoed, C. J. Terkelsen and J. F. Lassen *Circulation* (2008) 118; 1155-62

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18725489

Are "treatment" bare metal stents superior to "control" bare metal stents? A meta-analytic approach D. M. Kent and T. A. Trikalinos *Am Heart J* (2008) 155; 624-9, 29 e1-2

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18371468

Meta-analysis of the effects of paclitaxel-eluting stents versus bare metal stents on volumetric intravascular ultrasound in patients with versus without diabetes mellitus M. Kimura, G. S. Mintz, N. J. Weissman, K. D. Dawkins, E. Grube, S. G. Ellis, L. A. Cannon, Z. Masud, L. Mandinov, D. Baim and G. W. Stone *Am J Cardiol* (2008) 101; 1263-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18435955

Long-term outcomes of coronary-artery bypass graft surgery versus percutaneous coronary intervention for multivessel coronary artery disease in the bare-metal stent era T. Kimura, T. Morimoto, Y. Furukawa, Y. Nakagawa, S. Shizuta, N. Ehara, R. Taniguchi, T. Doi, K. Nishiyama, N. Ozasa, N. Saito, K. Hoshino, H. Mitsuoka, M. Abe, M. Toma, T. Tamura, Y. Haruna, Y. Imai, S. Teramukai, M. Fukushima and T. Kita *Circulation* (2008) 118; S199-209

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18824755

Three-year survival following multivessel percutaneous coronary intervention with bare-metal or drug-eluting stents in unselected patients N. Kukreja, Y. Onuma, H. M. Garcia-Garcia, J. Daemen, R. van Domburg and P. W. Serruys *Am J Cardiol* (2009) 103; 203-11

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19121437

Sirolimus-eluting stents, bare metal stents or coronary artery bypass grafting for patients with multivessel disease including involvement of the proximal left anterior descending artery: analysis of the Arterial Revascularization Therapies study part 2 (ARTS-II) N. Kukreja, P. W. Serruys, B. de Bruyne, A. Colombo, C. Macaya, G. Richardt,

J. Fajadet, C. Hamm, D. Goedhart, N. Macours and H. P. Stoll *Heart* (2009)
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19304671

An integrated TAXUS IV, V, and VI intravascular ultrasound analysis of the predictors of edge restenosis after bare metal or paclitaxel-eluting stents J. Liu, A. Maehara, G. S. Mintz, N. J. Weissman, A. Yu, H. Wang, L. Mandinov, J. J. Popma, S. G. Ellis, E. Grube, K. D. Dawkins and G. W. Stone *Am J Cardiol* (2009) 103; 501-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195510

Usefulness of serum cathepsin L as an independent biomarker in patients with coronary heart disease Y. Liu, X. Li, D. Peng, Z. Tan, H. Liu, Y. Qing, Y. Xue and G. P. Shi *Am J Cardiol* (2009) 103; 476-81
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195505

Comparison of stent thrombosis, myocardial infarction, and mortality following drug-eluting versus bare-metal stent coronary intervention in patients with diabetes mellitus M. Maeng, L. O. Jensen, A. Kaltoft, H. H. Hansen, M. Bottcher, J. F. Lassen, P. Thayssen, L. R. Krusell, K. Rasmussen, L. Pedersen, H. T. Sorensen, S. P. Johnsen and L. Thuesen *Am J Cardiol* (2008) 102; 165-72
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18602515

Comparison of effectiveness and safety of sirolimus-eluting stents versus bare-metal stents in patients with diabetes mellitus (from the Italian Multicenter Randomized DESSERT Study) A. Maresta, E. Varani, M. Balducelli, F. Varbella, C. Lettieri, L. Uguccioni, P. Sangiorgio and G. B. Zoccai *Am J Cardiol* (2008) 101; 1560-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18489933

Long-term clinical outcomes after drug-eluting and bare-metal stenting in Massachusetts L. Mauri, T. S. Silbaugh, R. E. Wolf, K. Zelevinsky, A. Lovett, Z. Zhou, F. S. Resnic and S. L. Normand *Circulation* (2008) 118; 1817-27
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18852368

Optimal treatment for in-stent restenosis after BMS--DES, coated balloon, or scalpel? D. Mukherjee *Eur Heart J* (2008) 29; 1595-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18556717

Drug-eluting stents versus bare metal stents for narrowing in saphenous vein grafts T. Okabe, J. Lindsay, A. N. Buch, D. H. Steinberg, P. Roy, T. L. Slottow, K. Smith, R. Torguson, Z. Xue, L. F. Satler, K. M. Kent, A. D. Pichard, N. J. Weissman and R. Waksman *Am J Cardiol* (2008) 102; 530-4
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18721507

Two-year clinical outcome with drug-eluting stents versus bare-metal stents in a real-world registry of unprotected left main coronary artery stenosis from the Italian Society of Invasive Cardiology T. Palmerini, A. Marzocchi, C. Tamburino, I. Sheiban, M. Margheri, G. Vecchi, G. Sangiorgi, A. Santarelli, A. Bartorelli, C. Briguori, L. Vignali, F.

Di Pede, A. Ramondo, L. Inglese, M. De Carlo, L. Bolognese, A. Benassi, C. Palmieri, V. Filippone, D. Sangiorgi and S. De Servi *Am J Cardiol* (2008) 102; 1463-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19026296

Comparison of six-month outcomes for primary percutaneous revascularization for acute myocardial infarction with drug-eluting versus bare metal stents (from the APEX-AMI study) M. R. Patel, M. E. Pfisterer, A. Betriu, P. Widmisky, D. R. Holmes, Jr., W. W. O'Neill, A. Stebbins, F. Van de Werf, P. W. Armstrong and C. B. Granger *Am J Cardiol* (2009) 103; 181-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19121433

Meta-analysis comparison (nine trials) of outcomes with drug-eluting stents versus bare metal stents in patients with diabetes mellitus G. Patti, A. Nusca and G. Di Sciascio *Am J Cardiol* (2008) 102; 1328-34

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18993150

Drug-eluting or bare-metal stents for large coronary vessel stenting? The BASKET-PROVE (PROspective Validation Examination) trial: study protocol and design M. Pfisterer, O. Bertel, P. O. Bonetti, H. P. Brunner-La Rocca, F. R. Eberli, P. Erne, S. Galatius, B. Hornig, W. Kiowski, O. Pachinger, G. Pedrazzini, H. Rickli, S. De Servi and C. Kaiser *Am Heart J* (2008) 155; 609-14

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18371466

The balance of risks and benefits of drug-eluting versus bare-metal stents M. Pfisterer, H. P. Brunner-La Rocca and C. Kaiser *J Am Coll Cardiol* (2008) 51; 972; author reply 72-3

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18308171

Long-term benefit-risk balance of drug-eluting vs. bare-metal stents in daily practice: does stent diameter matter? Three-year follow-up of BASKET M. Pfisterer, H. P. Brunner-La Rocca, P. Rickenbacher, P. Hunziker, C. Mueller, F. Nietlispach, G. Leibundgut, F. Bader and C. Kaiser *Eur Heart J* (2009) 30; 16-24

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19033260

Temporal changes in coronary revascularization procedures, outcomes, and costs in the bare-metal stent and drug-eluting stent eras: results from the US Medicare program J. Ryan, W. Linde-Zwirble, L. Engelhart, L. Cooper and D. J. Cohen *Circulation* (2009) 119; 952-61

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19204307

Long-term impact of drug-eluting stents versus bare-metal stents on all-cause mortality M. H. Shishehbor, S. S. Goel, S. R. Kapadia, D. L. Bhatt, P. Kelly, R. E. Raymond, J. M. Galla, S. J. Brener, P. L. Whitlow and S. G. Ellis *J Am Coll Cardiol* (2008) 52; 1041-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18848135

Mortality following placement of drug-eluting and bare-metal stents for ST-segment

elevation acute myocardial infarction in the Global Registry of Acute Coronary Events P. G. Steg, K. A. Fox, K. A. Eagle, M. Furman, F. Van de Werf, G. Montalescot, S. G. Goodman, A. Avezum, W. Huang and J. M. Gore *Eur Heart J* (2009) 30; 321-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19147604

Outcome differences with the use of drug-eluting stents for the treatment of in-stent restenosis of bare-metal stents versus drug-eluting stents D. H. Steinberg, M. A. Gaglia, Jr., T. L. Pinto Slottow, P. Roy, L. Bonello, A. De Labriolle, G. Lemesle, R. Torguson, K. Kineshige, Z. Xue, W. O. Suddath, K. M. Kent, L. F. Satler, A. D. Pichard, J. Lindsay and R. Waksman *Am J Cardiol* (2009) 103; 491-5
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195508

Are drug-eluting stents superior to bare-metal stents in patients with unprotected non-bifurcational left main disease? Insights from a multicentre registry C. Tamburino, M. E. Di Salvo, D. Capodanno, A. Marzocchi, I. Sheiban, M. Margheri, A. Maresta, F. Barlocco, G. Sangiorgi, G. Piovaccari, A. Bartorelli, C. Briguori, D. Ardissino, F. Di Pede, A. Ramondo, L. Inglese, A. S. Petronio, L. Bolognese, A. Benassi, C. Palmieri, A. Patti and S. De Servi *Eur Heart J* (2009)
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19276194

Comparison of drug-eluting stents and bare-metal stents for the treatment of unprotected left main coronary artery disease in acute coronary syndromes C. Tamburino, M. E. Di Salvo, D. Capodanno, T. Palmerini, I. Sheiban, M. Margheri, G. Vecchi, G. Sangiorgi, G. Piovaccari, A. Bartorelli, C. Briguori, D. Ardissino, F. Di Pede, A. Ramondo, L. Inglese, A. S. Petronio, L. Bolognese, A. Benassi, C. Palmieri, V. Filippone and S. De Servi *Am J Cardiol* (2009) 103; 187-93
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19121434

Cytochrome P450 2C19 681G>A polymorphism and high on-clopidogrel platelet reactivity associated with adverse 1-year clinical outcome of elective percutaneous coronary intervention with drug-eluting or bare-metal stents D. Trenk, W. Hochholzer, M. F. Fromm, L. E. Chialda, A. Pahl, C. M. Valina, C. Stratz, P. Schmiebusch, H. P. Bestehorn, H. J. Buttner and F. J. Neumann *J Am Coll Cardiol* (2008) 51; 1925-34
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18482659

Sirolimus-eluting stents versus bare-metal stents in patients with ST-segment elevation myocardial infarction: 9-month angiographic and intravascular ultrasound results and 12-month clinical outcome results from the MISSION! Intervention Study B. L. van der Hoeven, S. S. Liem, J. W. Jukema, N. Suraphakdee, H. Putter, J. Dijkstra, D. E. Atsma, M. Bootsma, K. Zeppenfeld, P. V. Oemrawsingh, E. E. van der Wall and M. J. Schalij *J Am Coll Cardiol* (2008) 51; 618-26
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18261680

Long-term outcomes with drug-eluting stents versus bare metal stents in the treatment of saphenous vein graft disease (results from the REgistro Regionale AngiopLastiche Emilia-Romagna registry) L. Vignali, F. Saia, A. Manari, A. Santarelli, A. Rubboli, E.

Varani, G. Piovaccari, A. Menozzi, G. Percoco, A. Benassi, G. Rusticali, P. Marzaroli, P. Guastaroba, R. Grilli, A. Maresta and A. Marzocchi *Am J Cardiol* (2008) 101; 947-52
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18359313

Survival advantage in Medicare patients receiving drug-eluting stents compared with bare metal stents: real or artefactual? F. W. Wang, B. F. Uretsky, J. L. Freeman, D. Zhang, S. H. Giordano and J. S. Goodwin *Catheter Cardiovasc Interv* (2008) 71; 636-43
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18360856

Long-term clinical outcomes after sirolimus-eluting stent implantation for treatment of restenosis within bare-metal versus drug-eluting stents C. Whan Lee, S. H. Kim, J. Suh, D. W. Park, S. H. Lee, Y. H. Kim, M. K. Hong, J. J. Kim, S. W. Park and S. J. Park *Catheter Cardiovasc Interv* (2008) 71; 594-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18311841