

1. *Extracorporeal cardiac shock wave therapy ameliorates clinical symptoms and improves regional myocardial blood flow in a patient with severe coronary artery disease and refractory angina*
Prinz, C., et al.
Case Report Med, 2009. **2009**: p. 639594.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19724656

2. *Reply to Sergej V. Jargin: Shock wave therapy of ischemic heart disease in the light of general pathology*
Holfeld, J., et al.
Int J Cardiol, 2009.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19758716

3. *Shock wave therapy of ischemic heart disease: Some aspects of publication and advertizing in Russia*
Jargin, S.V.
Int J Cardiol, 2009.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19910064

4. *Direct epicardial shock wave therapy improves ventricular function and induces angiogenesis in ischemic heart failure*
Zimpfer, D., et al.
J Thorac Cardiovasc Surg, 2009. **137**(4): p. 963-70.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19327525

5. *Extracorporeal shock wave therapy as a new and non-invasive angiogenic strategy*
Ito, K., Y. Fukumoto, and H. Shimokawa
Tohoku J Exp Med, 2009. **219**(1): p. 1-9.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19713678