

Effectiveness of "Direct" Stenting Without Balloon Predilatation (from the Multilink Tetra Randomised European Direct Stent Study [TRENDS]) K. D. Dawkins, B. Chevalier, M. J. Suttorp, L. Thuesen, E. Benit, A. Bethencourt, U. Morjaria, S. Veldhof, C. Dorange and A. van Weert

Am J Cardiol (2006) 97;316-21

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16442388](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16442388)

Comparison of C-reactive protein levels after coronary stenting with bare metal versus sirolimus-eluting stents J. M. de la Torre-Hernandez, F. Sainz-Laso, V. Burgos, T. Perez, A. Figueroa, J. Zueco and T. Colman

Am J Cardiol (2005) 95;748-51

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15757602](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15757602)

Influence of stent surface topography on the outcomes of patients undergoing coronary stenting: a randomized double-blind controlled trial A. Dibra, A. Kastrati, J. Mehilli, J.

Pache, R. von Oepen, J. Dirschinger and A. Schomig

Catheter Cardiovasc Interv (2005) 65;374-80

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15926182](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15926182)

Comparison of C-reactive protein levels before and after coronary stenting and restenosis among patients treated with sirolimus-eluting versus bare metal stents A. Dibra, G. Ndreppepa, J. Mehilli, J. Dirschinger, J. Pache, H. Schuhlen, A. Schomig and A. Kastrati

Am J Cardiol (2005) 95;1238-40

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15878000](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15878000)

Two-year follow-up intravascular ultrasound analysis after bare metal stent implantation in 120 lesions M. K. Hong, C. W. Lee, Y. H. Kim, B. K. Lee, M. K. Kim, T. H. Yang, J. M. Song, K. H. Han, D. H. Kang, J. K. Song, J. J. Kim, S. W. Park and S. J. Park

Catheter Cardiovasc Interv (2005) 65;247-53

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15858789](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15858789)

Drug-eluting stents versus bare metal stents in percutaneous coronary interventions (a meta-analysis) C. Indolfi, M. Pavia and I. F. Angelillo

Am J Cardiol (2005) 95;1146-52

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15877984](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15877984)

Effect of lumen narrowing within coronary stents on proximal and distal vessel segments following bare metal stent implantation H. Kaneda, J. Ako, T. Kataoka, A. Miyazawa, M. Terashima, F. Ikeno, S. Sonoda, Y. Shimada, Y. Morino, Y. Honda, P. G. Yock and P. J. Fitzgerald

Am J Cardiol (2005) 96;376-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16054461](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16054461)

Meta-analysis comparing drug-eluting stents with bare metal stents D. G. Katritsis, E. Karvouni and J. P. Ioannidis

Am J Cardiol (2005) 95;640-3

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15721109](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15721109)

Comparison of effects of drug-eluting stents versus bare metal stents on plasma C-reactive protein levels J. Y. Kim, Y. G. Ko, C. Y. Shim, S. Park, K. C. Hwang, D. Choi, Y. Jang, N. Chung, W. H. Shim and S. Y. Cho

Am J Cardiol (2005) 96;1384-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16275183](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16275183)

Randomized comparison of carbon ion-implanted stent versus bare metal stent in coronary artery disease: the Asian Pacific Multicenter Arthos Stent Study (PASS) trial Y. H. Kim, C. W. Lee, M. K. Hong, S. W. Park, S. J. Tahk, J. Y. Yang, S. Saito, T. Santoso, L. Quan, J. Ge, N. J. Weissman, A. J. Lansky, G. S. Mintz and S. J. Park

Am Heart J (2005) 149;336-41

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15846274](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15846274)

Comparison of sirolimus-eluting stent, paclitaxel-eluting stent, and bare metal stent in the treatment of long coronary lesions Y. H. Kim, S. W. Park, C. W. Lee, M. K. Hong, H. C. Gwon, Y. Jang, M. M. Lee, B. K. Koo, D. J. Oh, K. B. Seung, S. J. Tahk, J. Yoon and S. J. Park

Catheter Cardiovasc Interv (2006) 67;181-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16400663](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16400663)

Clinical and angiographic analysis with a cobalt alloy coronary stent (driver) in stable and unstable angina pectoris V. Legrand, H. Kelbaek, K. E. Hauptmann, D. Glogar, W. Rutsch, G. Grollier, P. Vermeersch, J. Elias and C. C. De Cock

Am J Cardiol (2006) 97;349-52

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16442394](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16442394)

Comparison of effectiveness of sirolimus-eluting stents versus bare metal stents for percutaneous coronary intervention in patients at high risk for coronary restenosis or clinical adverse events A. Marzocchi, G. Piovaccari, A. Manari, E. Aurier, A. Benassi, F. Saia, G. Casella, E. Varani, A. Santarelli, P. Guastaroba, R. Grilli and A. Maresta

Am J Cardiol (2005) 95;1409-14

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15950561](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15950561)

Comparison of thrombosis and restenosis risk from stent length of sirolimus-eluting stents versus bare metal stents L. Mauri, A. J. O'Malley, J. J. Popma, J. W. Moses, M. B. Leon, D. R. Holmes, Jr., P. S. Teirstein, D. E. Cutlip, D. Donahoe and R. E. Kuntz

Am J Cardiol (2005) 95;1140-5

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15877983](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15877983)

Comparison of quantitative angiographic parameters with the magnitude of neointimal hyperplasia measured by volumetric intravascular ultrasound in patients treated with bare metal and nonpolymeric paclitaxel-coated stents G. S. Mintz, M. K. Hong, A. E. Raizner, C. W. Lee, J. J. Kim, E. Escolar, N. E. Fearnott, S. W. Park, S. J. Park and N. J.

- Weissman  
Am J Cardiol (2005) 95;105-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15619403](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15619403)  
Drug-eluting stents compared with thin-strut bare stents for the reduction of restenosis: a prospective, randomized trial J. Pache, A. Dibra, J. Mehilli, J. Dirschinger, A. Schomig and A. Kastrati
- Eur Heart J (2005) 26;1262-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15737962](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15737962)  
Safety of magnetic resonance imaging one to three days after bare metal and drug-eluting stent implantation I. Porto, J. Selvanayagam, V. Ashar, S. Neubauer and A. P. Banning
- Am J Cardiol (2005) 96;366-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16054459](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16054459)  
Primary stenting of occluded native coronary arteries II--rationale and design of the PRISON II study: a randomized comparison of bare metal stent implantation with sirolimus-eluting stent implantation for the treatment of chronic total coronary occlusions B. M. Rahel, G. J. Laarman and M. J. Suttorp
- Am Heart J (2005) 149;e1-3  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15864219](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15864219)  
Randomized comparison of sirolimus-eluting stent versus standard stent for percutaneous coronary revascularization in diabetic patients: the diabetes and sirolimus-eluting stent (DIABETES) trial M. Sabate, P. Jimenez-Quevedo, D. J. Angiolillo, J. A. Gomez-Hospital, F. Alfonso, R. Hernandez-Antolin, J. Goicolea, C. Banuelos, J. Escaned, R. Moreno, C. Fernandez, F. Fernandez-Aviles and C. Macaya Circulation (2005) 112;2175-83  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16203930](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16203930)  
Prospective randomized comparison of early and late results of a carbonized stent versus a high-grade stainless steel stent of identical design: the PREVENT Trial [corrected] P. B. Sick, O. Brosteanu, M. Ulrich, H. Thiele, J. Niebauer, I. Busch and G. Schuler
- Am Heart J (2005) 149;681-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15990753](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15990753)  
Evaluation of the Medtronic (Driver) cobalt-chromium alloy coronary stent system M. H. Sketch, Jr., M. Ball, B. Rutherford, J. J. Popma, C. Russell and D. J. Kereiakes
- Am J Cardiol (2005) 95;8-12  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15619386](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15619386)  
Circadian variation in symptomatic subacute stent thrombosis after bare metal coronary stent implantation A. Tamura, T. Watanabe, K. Nagase, T. Nakaishi, N. Aso, Y. Kawano, Y. Abe, H. Zaizen, S. Yano and J. Kadota

Am J Cardiol (2006) 97;195-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16442362](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16442362)

Comparison of bare-metal stenting with minimally invasive bypass surgery for stenosis of the left anterior descending coronary artery: a 5-year follow-up H. Thiele, S. Oettel, S. Jacobs, R. Hambrecht, P. Sick, J. F. Gummert, F. W. Mohr, G. Schuler and V. Falk  
Circulation (2005) 112;3445-50

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16316966](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16316966)

Impact and temporal trends of percutaneous coronary intervention in the drug-eluting stent versus bare metal stent eras C. A. Thompson, A. V. Kaplan, J. E. Jayne, B. J. Friedman, B. D. Hettelman, N. W. Niles and J. F. Robb

Am J Cardiol (2005) 96;668-72

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=16125492](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16125492)

Stent thrombosis following bare-metal stent implantation: success of emergency percutaneous coronary intervention and predictors of adverse outcome P. Wenaweser, C. Rey, F. R. Eberli, M. Togni, D. Tuller, S. Locher, A. Remondino, C. Seiler, O. M. Hess, B. Meier and S. Windecker

Eur Heart J (2005) 26;1180-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15728650](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15728650)

Angiographic, intravascular ultrasound, and fractional flow reserve evaluation of direct stenting vs. conventional stenting using BeStent2 in a multicentre randomized trial W. Wijns, S. Verheye, G. Manoharan, G. S. Werner, E. Grube, B. De Bruyne, J. Koolen, C. W. Hamm, A. Medina, J. W. Bech and P. J. De Feyter

Eur Heart J (2005) 26;1852-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15888499](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15888499)

Influence of stent length to lesion length ratio on angiographic and clinical outcomes after implantation of bare metal and drug-eluting stents (the TAXUS-IV Study) S. C. Wong, M. K. Hong, S. G. Ellis, M. Buchbinder, M. Reisman, A. DeLago, M. Kellett, J. J. Popma, M. E. Russell, R. Mehran, J. W. Moses and G. W. Stone

Am J Cardiol (2005) 95;1043-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15842968](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15842968)

Effect of bare metal stenting on angiographic and clinical outcomes in diabetic and nondiabetic patients undergoing percutaneous coronary intervention of nonacute occluded coronary arteries: a report from the total occlusion study of Canada (TOSCA) K. M. Yee, C. E. Buller, D. Catellier, E. A. Cohen, R. C. Carere, T. Anderson, P. Berger, J. R. Burton, G. Barbeau, K. K. Teo and V. Dzavik

Catheter Cardiovasc Interv (2005) 66;178-84

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=15977265](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15977265)