

Frequency of myocardial bridges and dynamic compression of epicardial coronary arteries: a comparison between computed tomography and invasive coronary angiography P. J. Kim, G. Hur, S. Y. Kim, J. Namgung, S. W. Hong, Y. H. Kim and W. R. Lee *Circulation* (2009) 119; 1408-16

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19255347

Prognostic value of normal adenosine-stress cardiac magnetic resonance imaging G. Pilz, A. Jeske, M. Klos, E. Ali, B. Hoefling, R. Scheck and P. Bernhardt *Am J Cardiol* (2008) 101; 1408-12

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18471450

Simultaneous "hybrid" percutaneous coronary intervention and minimally invasive surgical bypass grafting: feasibility, safety, and clinical outcomes B. Reicher, R. S. Poston, M. R. Mehra, A. Joshi, P. Odonkor, Z. Kon, P. A. Reyes and D. A. Zimrin *Am Heart J* (2008) 155; 661-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18371473