

Hyperhomocysteinemia, a cardiac metabolic disease: role of nitric oxide and the p22phox subunit of NADPH oxidase J. S. Becker, A. Adler, A. Schneeberger, H. Huang, Z. Wang, E. Walsh, A. Koller and T. H. Hintze

Circulation (2005) 111;2112-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15851618

Comparison of usefulness of inflammatory markers in patients with versus without peripheral arterial disease in predicting adverse cardiovascular outcomes (myocardial infarction, stroke, and death) J. A. Beckman, O. Preis, P. M. Ridker and M.

Gerhard-Herman

Am J Cardiol (2005) 96;1374-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16275181

Monocyte cyclooxygenase-2 overactivity: a new marker of subclinical atherosclerosis in asymptomatic subjects with cardiovascular risk factors? O. Beloqui, J. A. Paramo, J.

Orbe, A. Benito, I. Colina, A. Monasterio and J. Diez

Eur Heart J (2005) 26;153-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15618071

C-reactive protein as a predictor of cardiovascular risk in a population with a high prevalence of diabetes: the Strong Heart Study L. G. Best, Y. Zhang, E. T. Lee, J. L. Yeh, L. Cowan, V. Palmieri, M. Roman, R. B. Devereux, R. R. Fabsitz, R. P. Tracy, D. Robbins, M. Davidson, A. Ahmed and B. V. Howard

Circulation (2005) 112;1289-95

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16116058

Efficacy and safety of high-density lipoprotein cholesterol-increasing compounds: a meta-analysis of randomized controlled trials R. S. Birjmohun, B. A. Hutten, J. J.

Kastelein and E. S. Stroes

J Am Coll Cardiol (2005) 45;185-97

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15653014

The relationship between C-reactive protein and subclinical cardiovascular disease in the Diabetes Heart Study (DHS) D. W. Bowden, L. A. Lange, C. D. Langefeld, K. B.

Brosnihan, B. I. Freedman, J. J. Carr, L. E. Wagenknecht and D. M. Herrington

Am Heart J (2005) 150;1032-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16290992

Association of lipoprotein-associated phospholipase A2 levels with coronary artery disease risk factors, angiographic coronary artery disease, and major adverse events at follow-up E. S. Brilakis, J. P. McConnell, R. J. Lennon, A. A. Elesber, J. G. Meyer and P. B. Berger

Eur Heart J (2005) 26;137-44

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15618069

Impact of the elevation of biochemical markers of myocardial damage on long-term

mortality after percutaneous coronary intervention: results of the CK-MB and PCI study
C. Cavallini, S. Savonitto, R. Violini, G. Arraiz, M. Plebani, Z. Olivari, P. Rubartelli, S. Battaglia, L. Niccoli, G. Steffenino and D. Ardissino
Eur Heart J (2005) 26;1494-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15741227

Near-infrared fluorescent imaging of matrix metalloproteinase activity after myocardial infarction J. Chen, C. H. Tung, J. R. Allport, S. Chen, R. Weissleder and P. L. Huang
Circulation (2005) 111;1800-5

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15809374

Comparison of differing C-reactive protein assay methods and their impact on cardiovascular risk assessment J. L. Clarke, J. L. Anderson, J. F. Carlquist, R. F. Roberts, B. D. Horne, T. L. Bair, M. J. Kolek, C. P. Mower, A. M. Crane, W. L. Roberts and J. B. Muhlestein
Am J Cardiol (2005) 95;155-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15619419

Relationship among pregnancy associated plasma protein-A levels, clinical characteristics, and coronary artery disease extent in patients with chronic stable angina pectoris J. Cosin-Sales, J. C. Kaski, M. Christiansen, P. Kaminski, C. Oxvig, M. T. Overgaard, D. Cole and D. W. Holt
Eur Heart J (2005) 26;2093-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16055491

C-reactive protein and the 10-year incidence of coronary heart disease in older men and women: the cardiovascular health study M. Cushman, A. M. Arnold, B. M. Psaty, T. A. Manolio, L. H. Kuller, G. L. Burke, J. F. Polak and R. P. Tracy
Circulation (2005) 112;25-31

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15983251

Comparison of C-reactive protein levels after coronary stenting with bare metal versus sirolimus-eluting stents J. M. de la Torre-Hernandez, F. Sainz-Laso, V. Burgos, T. Perez, A. Figueroa, J. Zueco and T. Colman
Am J Cardiol (2005) 95;748-51

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15757602

Inflammatory response after intervention assessed by serial C-reactive protein measurements correlates with restenosis in patients treated with coronary stenting A. Dibra, J. Mehilli, S. Braun, M. Hadamitzky, H. Baum, J. Dirschinger, H. Schuhlen, A. Schomig and A. Kastrati
Am Heart J (2005) 150;344-50

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16086941

Comparison of C-reactive protein levels before and after coronary stenting and restenosis among patients treated with sirolimus-eluting versus bare metal stents A.

Dibra, G. Ndrepepa, J. Mehilli, J. Dirschinger, J. Pache, H. Schuhlen, A. Schomig and A. Kastrati

Am J Cardiol (2005) 95;1238-40

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15878000

Possible role of brain-derived neurotrophic factor in the pathogenesis of coronary artery disease J. Ejiri, N. Inoue, S. Kobayashi, R. Shiraki, K. Otsui, T. Honjo, M. Takahashi, Y. Ohashi, S. Ichikawa, M. Terashima, T. Mori, K. Awano, T. Shinke, J. Shite, K. Hirata, H. Yokozaki, S. Kawashima and M. Yokoyama

Circulation (2005) 112;2114-20

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16186425

Urinary albumin excretion is independently associated with carotid and femoral artery atherosclerosis in the general population M. Furtner, S. Kiechl, A. Mair, K. Seppi, S. Weger, F. Oberholzer, W. Poewe and J. Willeit

Eur Heart J (2005) 26;279-87

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15618058

Serum soluble lectin-like oxidized low-density lipoprotein receptor-1 levels are elevated in acute coronary syndrome: a novel marker for early diagnosis K. Hayashida, N. Kume, T. Murase, M. Minami, D. Nakagawa, T. Inada, M. Tanaka, A. Ueda, G. Kominami, H. Kambara, T. Kimura and T. Kita

Circulation (2005) 112;812-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16061745

Pregnancy-associated plasma protein-A levels in patients with acute coronary syndromes: comparison with markers of systemic inflammation, platelet activation, and myocardial necrosis C. Heeschen, S. Dimmeler, C. W. Hamm, S. Fichtlscherer, M. L. Simoons and A. M. Zeiher

J Am Coll Cardiol (2005) 45;229-37

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15653020

Increased circulating C-reactive protein and macrophage-colony stimulating factor are complementary predictors of long-term outcome in patients with chronic coronary artery disease I. Ikonomidis, J. Lekakis, I. Revela, F. Andreotti and P. Nihoyannopoulos

Eur Heart J (2005) 26;1618-24

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15800017

Local release of C-reactive protein from vulnerable plaque or coronary arterial wall injured by stenting T. Inoue, T. Kato, T. Uchida, M. Sakuma, A. Nakajima, M. Shibasaki, Y. Imoto, M. Saito, S. Hashimoto, Y. Hikichi and K. Node

J Am Coll Cardiol (2005) 46;239-45

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16022949

Usefulness of biomarkers for predicting long-term mortality in patients with diabetes mellitus and non-ST-elevation acute coronary syndromes (a GUSTO IV substudy) S. K.

- James, B. Lindahl, J. R. Timmer, J. P. Ottervanger, A. Siegbahn, M. Stridsberg, P. Armstrong, R. Califf, L. Wallentin and M. L. Simoons
Am J Cardiol (2006) 97;167-72
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16442356
- Relationship between C-reactive protein and subclinical atherosclerosis: the Dallas Heart Study A. Khera, J. A. de Lemos, R. M. Peshock, H. S. Lo, H. G. Stanek, S. A. Murphy, F. H. Wians, Jr., S. M. Grundy and D. K. McGuire
Circulation (2006) 113;38-43
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16380546
- Race and gender differences in C-reactive protein levels A. Khera, D. K. McGuire, S. A. Murphy, H. G. Stanek, S. R. Das, W. Vongpatanasin, F. H. Wians, Jr., S. M. Grundy and J. A. de Lemos
J Am Coll Cardiol (2005) 46;464-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16053959
- Serum blood urea nitrogen as an independent marker of subsequent mortality among patients with acute coronary syndromes and normal to mildly reduced glomerular filtration rates A. J. Kirtane, D. M. Leder, S. S. Waikar, G. M. Chertow, K. K. Ray, D. S. Pinto, D. Karmpaliotis, A. J. Burger, S. A. Murphy, C. P. Cannon, E. Braunwald and C. M. Gibson
J Am Coll Cardiol (2005) 45;1781-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15936606
- Inflammatory markers and the metabolic syndrome: insights from therapeutic interventions K. K. Koh, S. H. Han and M. J. Quon
J Am Coll Cardiol (2005) 46;1978-85
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16325028
- The relationship between blood pressure and C-reactive protein in the Multi-Ethnic Study of Atherosclerosis (MESA) S. G. Lakoski, M. Cushman, W. Palmas, R. Blumenthal, R. B. D'Agostino, Jr. and D. M. Herrington
J Am Coll Cardiol (2005) 46;1869-74
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16286174
- Elevated placental growth factor levels are associated with adverse outcomes at four-year follow-up in patients with acute coronary syndromes T. Lenderink, C. Heeschen, S. Fichtlscherer, S. Dimmeler, C. W. Hamm, A. M. Zeiher, M. L. Simoons and E. Boersma
J Am Coll Cardiol (2006) 47;307-11
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16412852
- Serial analyses of N-terminal pro-B-type natriuretic peptide in patients with non-ST-segment elevation acute coronary syndromes: a Fragmin and fast Revascularisation during In Stability in Coronary artery disease (FRISC)-II substudy B.

- Lindahl, J. Lindback, T. Jernberg, N. Johnston, M. Stridsberg, P. Venge and L. Wallentin
J Am Coll Cardiol (2005) 45;533-41
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15708700
Low plasma adiponectin levels predict progression of coronary artery calcification D. M. Maahs, L. G. Ogden, G. L. Kinney, P. Wadwa, J. K. Snell-Bergeon, D. Dabelea, J. E. Hokanson, J. Ehrlich, R. H. Eckel and M. Rewers
Circulation (2005) 111;747-53
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15699257
Inflammatory markers at the site of ruptured plaque in acute myocardial infarction: locally increased interleukin-6 and serum amyloid A but decreased C-reactive protein W. Maier, L. A. Altwegg, R. Corti, S. Gay, M. Hersberger, F. E. Maly, G. Sutsch, M. Roffi, M. Neidhart, F. R. Eberli, F. C. Tanner, S. Gobbi, A. von Eckardstein and T. F. Luscher
Circulation (2005) 111;1355-61
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15753219
Circulating secretory phospholipase A2 activity predicts recurrent events in patients with severe acute coronary syndromes Z. Mallat, P. G. Steg, J. Benessiano, M. L. Tanguy, K. A. Fox, J. P. Collet, O. H. Dabbous, P. Henry, K. F. Carruthers, A. Dauphin, C. S. Arguelles, J. Masliah, B. Hugel, G. Montalescot, J. M. Freyssinet, B. Asselain and A. Tedgui
J Am Coll Cardiol (2005) 46;1249-57
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16198839
Diffuse and active inflammation occurs in both vulnerable and stable plaques of the entire coronary tree: a histopathologic study of patients dying of acute myocardial infarction A. Mauriello, G. Sangiorgi, S. Fratoni, G. Palmieri, E. Bonanno, L. Anemona, R. S. Schwartz and L. G. Spagnoli
J Am Coll Cardiol (2005) 45;1585-93
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15893171
Enhanced T-helper-1 lymphocyte activation patterns in acute coronary syndromes H. Methe, S. Brunner, D. Wiegand, M. Nabauer, J. Koglin and E. R. Edelman
J Am Coll Cardiol (2005) 45;1939-45
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15963390
C-reactive protein and cardiovascular outcomes in smokers versus nonsmokers in non-ST-elevation acute coronary syndrome (from the TACTICS-TIMI 18 trial) N. Moghbali, A. J. Kirtane, K. K. Ray, S. A. Murphy, C. M. Gibson, E. Braunwald and C. P. Cannon
Am J Cardiol (2005) 96;635-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16125484
Persistent systemic inflammation in unstable angina is largely unrelated to the

- atherothrombotic burden C. Monaco, E. Rossi, D. Milazzo, F. Citterio, F. Ginnetti, G. D'Onofrio, D. Cianflone, F. Crea, L. M. Biasucci and A. Maseri
J Am Coll Cardiol (2005) 45;238-43
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15653021
- Genetic inflammatory factors predict restenosis after percutaneous coronary interventions P. S. Monraats, N. M. Pires, W. R. Agema, A. H. Zwinderman, A. Schepers, M. P. de Maat, P. A. Doevedans, R. J. de Winter, R. A. Tio, J. Waltenberger, R. R. Frants, P. H. Quax, B. J. van Vlijmen, D. E. Atsma, A. van der Laarse, E. E. van der Wall and J. W. Jukema
Circulation (2005) 112;2417-25
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16230497
- Lipoprotein-associated phospholipase A2 activity is associated with risk of coronary heart disease and ischemic stroke: the Rotterdam Study H. H. Oei, I. M. van der Meer, A. Hofman, P. J. Koudstaal, T. Stijnen, M. M. Breteler and J. C. Witteman
Circulation (2005) 111;570-5
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15699277
- C-reactive protein and electrocardiographic ST-segment depression additively predict mortality: the Strong Heart Study P. M. Okin, M. J. Roman, L. G. Best, E. T. Lee, J. M. Galloway, B. V. Howard and R. B. Devereux
J Am Coll Cardiol (2005) 45;1787-93
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15936607
- Preprocedural levels of C-reactive protein and leukocyte counts predict 9-month mortality after coronary angioplasty for the treatment of unprotected left main coronary artery stenosis T. Palmerini, A. Marzocchi, C. Marrozzini, P. Ortolani, F. Saia, L. Bacchi-Reggiani, S. Virzi, S. Gianstefani and A. Branzi
Circulation (2005) 112;2332-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16203907
- Predictors of the rise in vWF after ST elevation myocardial infarction: implications for treatment strategies and clinical outcome: An ENTIRE-TIMI 23 substudy K. K. Ray, D. A. Morrow, C. M. Gibson, S. Murphy, E. M. Antman and E. Braunwald
Eur Heart J (2005) 26;440-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15673542
- Resistin is an inflammatory marker of atherosclerosis in humans M. P. Reilly, M. Lehrke, M. L. Wolfe, A. Rohatgi, M. A. Lazar and D. J. Rader
Circulation (2005) 111;932-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15710760
- Relative efficacy of atorvastatin 80 mg and pravastatin 40 mg in achieving the dual goals of low-density lipoprotein cholesterol <70 mg/dl and C-reactive protein <2 mg/l: an analysis of the PROVE-IT TIMI-22 trial P. M. Ridker, D. A. Morrow, L. M. Rose, N. Rifai,

- C. P. Cannon and E. Braunwald
J Am Coll Cardiol (2005) 45;1644-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15893181
Association of hemoglobin levels with clinical outcomes in acute coronary syndromes M. S. Sabatine, D. A. Morrow, R. P. Giugliano, P. B. Burton, S. A. Murphy, C. H. McCabe, C. M. Gibson and E. Braunwald
Circulation (2005) 111;2042-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15824203
Stent implantation, but not pathogen burden, is associated with plasma C-reactive protein and interleukin-6 levels after percutaneous coronary intervention in patients with stable angina pectoris N. Saleh, B. Svane, J. Jensen, L. O. Hansson, M. Nordin and P. Tornvall
Am Heart J (2005) 149;876-82
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15894971
CD40 ligand enhances monocyte tissue factor expression and thrombin generation via oxidative stress in patients with hypercholesterolemia V. Sanguigni, D. Ferro, P. Pignatelli, M. Del Ben, T. Nadia, M. Saliola, R. Sorge and F. Violi
J Am Coll Cardiol (2005) 45;35-42
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15629370
Glutathione peroxidase-1 and homocysteine for cardiovascular risk prediction: results from the AtheroGene study R. Schnabel, K. J. Lackner, H. J. Rupprecht, C. Espinola-Klein, M. Torzewski, E. Lubos, C. Bickel, F. Cambien, L. Tiret, T. Munzel and S. Blankenberg
J Am Coll Cardiol (2005) 45;1631-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15893179
Enhanced plasma levels of LIGHT in unstable angina: possible pathogenic role in foam cell formation and thrombosis H. Scholz, W. Sandberg, J. K. Damas, C. Smith, A. K. Andreassen, L. Gullestad, S. S. Froland, A. Yndestad, P. Aukrust and B. Halvorsen
Circulation (2005) 112;2121-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16186421
White blood cell count predicts reduction in coronary heart disease mortality with pravastatin R. A. Stewart, H. D. White, A. C. Kirby, S. R. Heritier, R. J. Simes, P. J. Nestel, M. J. West, D. M. Colquhoun and A. M. Tonkin
Circulation (2005) 111;1756-62
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15809366
Fasting glucose is an important independent risk factor for 30-day mortality in patients with acute myocardial infarction: a prospective study M. Suleiman, H. Hammerman, M. Boulos, M. R. Kapeliovich, A. Suleiman, Y. Agmon, W. Markiewicz and D. Aronson
Circulation (2005) 111;754-60

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15699267

Serum hepatocyte growth factor levels predict long-term clinical outcome after percutaneous coronary revascularization S. Susen, K. Sautiere, F. Mouquet, F. Cuilleret, A. Chmait, E. P. McFadden, B. Hennache, F. Richard, P. de Groote, J. M. Lablanche, J. Dallongeville, C. Bauters, B. Jude and E. Van Belle

Eur Heart J (2005) 26;2387-95

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16105849

Multiple plaque rupture and C-reactive protein in acute myocardial infarction A. Tanaka, K. Shimada, T. Sano, M. Namba, T. Sakamoto, Y. Nishida, T. Kawarabayashi, D. Fukuda and J. Yoshikawa

J Am Coll Cardiol (2005) 45;1594-9

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15893172

Elevated plaque temperature in non-culprit de novo atheromatous lesions of patients with acute coronary syndromes K. Toutouzas, M. Drakopoulou, J. Mitropoulos, E. Tsiamis, S. Vaina, M. Vavuranakis, V. Markou, E. Bosinakou and C. Stefanadis

J Am Coll Cardiol (2006) 47;301-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16412851

Plasma levels of soluble tumor necrosis factor receptor type I during the acute phase following complicated myocardial infarction predicts survival in high-risk patients T. Ueland, J. Kjekshus, S. S. Froland, T. Omland, I. B. Squire, L. Gullestad, K. Dickstein and P. Aukrust

J Am Coll Cardiol (2005) 46;2018-21

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=16325035

Associations between cigarette smoking, pipe/cigar smoking, and smoking cessation, and haemostatic and inflammatory markers for cardiovascular disease S. G.

Wannamethee, G. D. Lowe, A. G. Shaper, A. Rumley, L. Lennon and P. H. Whincup

Eur Heart J (2005) 26;1765-73

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15817606