

1. *Provisional TAP-stenting strategy to treat bifurcated lesions with drug-eluting stents: one-year clinical results of a prospective registry*  
Burzotta, F., et al.  
J Invasive Cardiol, 2009. **21**(10): p. 532-7.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19805841](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19805841)
  
2. *Randomized comparison of coronary bifurcation stenting with the crush versus the culotte technique using sirolimus eluting stents: the Nordic stent technique study*  
Erglis, A., et al.  
Circ Cardiovasc Interv, 2009. **2**(1): p. 27-34.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20031690](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20031690)
  
3. *Intervention for jailed diagonal branch is not necessary while stenting the left anterior descending artery*  
Ikeda, N., et al.  
J Interv Cardiol, 2009. **22**(3): p. 252-6.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19366401](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19366401)
  
4. *Double versus single stenting for coronary bifurcation lesions: a meta-analysis*  
Katritsis, D.G., G.C. Siontis, and J.P. Ioannidis  
Circ Cardiovasc Interv, 2009. **2**(5): p. 409-15.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20031750](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20031750)
  
5. *Physiologic evaluation of bifurcation lesions using fractional flow reserve*  
Koo, B.K.  
J Interv Cardiol, 2009. **22**(2): p. 110-3.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19379467](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19379467)
  
6. *Interventional therapy of bifurcation lesions: a TIMI flow-guided concept to treat side branches in bifurcation lesions--a prospective randomized clinical study (Thueringer bifurcation study, THUEBIS study as pilot trial)*  
Korn, H.V., et al.  
Circ Cardiovasc Interv, 2009. **2**(6): p. 535-42.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20031771](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20031771)

7. *Quantitative angiographic methods for bifurcation lesions: a consensus statement from the European Bifurcation Group*  
Lansky, A., et al.  
Catheter Cardiovasc Interv, 2009. **73**(2): p. 258-66.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19085918](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19085918)
  
8. *Ostial and midshaft lesions vs. bifurcation lesions in 1111 patients with unprotected left main coronary artery stenosis treated with drug-eluting stents: results of the survey from the Italian Society of Invasive Cardiology*  
Palmerini, T., et al.  
Eur Heart J, 2009. **30**(17): p. 2087-94.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19508996](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19508996)
  
9. *Ostial and midshaft lesions vs. bifurcation lesions in 1111 patients with unprotected left main coronary artery stenosis treated with drug-eluting stents: results of the survey from the Italian Society of Invasive Cardiology*  
Palmerini, T., et al.  
Eur Heart J, 2009. **30**(17): p. 2087-94.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19508996](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19508996)
  
10. *Evaluation of a bifurcation drug-eluting stent system versus provisional T-stenting in a perfused synthetic coronary artery model*  
Rizik, D.G., et al.  
J Interv Cardiol, 2009. **22**(6): p. 537-46.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19912465](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19912465)
  
11. *Very long-term results comparing a simple versus a complex stenting strategy in the treatment of coronary bifurcation lesions*  
Sheiban, I., et al.  
Catheter Cardiovasc Interv, 2009. **74**(2): p. 313-20.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19496110](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19496110)
  
12. *Three-year outcome of drug-eluting stent implantation for coronary artery bifurcation lesions*

Stinis, C.T., et al.

Catheter Cardiovasc Interv, 2009.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19924774](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19924774)

13. *Strategies for drug-eluting stent treatment of bifurcation coronary artery disease in the United States: insights from the e-Cypher S.T.L.L.R.trial*

Suzuki, N., et al.

Catheter Cardiovasc Interv, 2009. **73**(7): p. 890-7.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19455662](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19455662)

14. *9-month clinical, angiographic, and intravascular ultrasound results of a prospective evaluation of the Axxess self-expanding biolimus A9-eluting stent in coronary bifurcation lesions: the DIVERGE (Drug-Eluting Stent Intervention for Treating Side Branches Effectively) study*

Verheye, S., et al.

J Am Coll Cardiol, 2009. **53**(12): p. 1031-9.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19298915](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19298915)

15. *Long-term clinical and angiographic outcomes after implantation of sirolimus-eluting stents with a "modified mini-crush" technique in coronary bifurcation lesions*

Yang, H.M., et al.

Catheter Cardiovasc Interv, 2009. **74**(1): p. 76-84.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19360861](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19360861)

16. *Histone deacetylase 3 is critical in endothelial survival and atherosclerosis development in response to disturbed flow*

Zampetaki, A., et al.

Circulation, 2010. **121**(1): p. 132-42.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20026773](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20026773)

17. *Simple versus complex stenting strategy for coronary artery bifurcation lesions in the drug-eluting stent era: a meta-analysis of randomised trials*

Zhang, F., L. Dong, and J. Ge

Heart, 2009. **95**(20): p. 1676-81.

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list\_uids=19643768

18. *Coronary bifurcation lesions: to stent one branch or both? A meta-analysis of patients treated with drug eluting stents*

Niccoli, G., et al.

Int J Cardiol, 2010. **139**(1): p. 80-91.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19027969](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19027969)