

1. *Outcomes after percutaneous coronary intervention of ostial lesions in the era of drug-eluting stents*

Freeman, M., et al.

Catheter Cardiovasc Interv, 2009. **73**(6): p. 763-8.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19309731](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19309731)

2. *Ostial and midshaft lesions vs. bifurcation lesions in 1111 patients with unprotected left main coronary artery stenosis treated with drug-eluting stents: results of the survey from the Italian Society of Invasive Cardiology*

Palmerini, T., et al.

Eur Heart J, 2009. **30**(17): p. 2087-94.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19508996](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19508996)

3. *Emergency percutaneous coronary intervention in the setting of a single left coronary ostium*

Williams, P.D., R.W. Bury, and M.J. Brack

Circ Cardiovasc Interv, 2009. **2**(2): p. 149-51.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=20031709](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20031709)