

- Survival benefit after percutaneous treatment of chronic total coronary occlusions G. Belli Eur Heart J (2009) 30; 506; author reply 06-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19168525
- Potential impact of noncontrast computed tomography as gatekeeper for myocardial perfusion positron emission tomography in patients admitted to the chest pain unit F. P. Esteves, R. Sanyal, C. A. Santana, L. Shaw and P. Raggi Am J Cardiol (2008) 101; 149-52
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18178397
- Non-invasive detection and quantification of acute myocardial infarction in rabbits using mono-[123I]iodohypericin microSPECT H. Fonge, K. Vunckx, H. Wang, Y. Feng, L. Mortelmans, J. Nuysts, G. Bormans, A. Verbruggen and Y. Ni Eur Heart J (2008) 29; 260-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18156139
- Interaction of caffeine with regadenoson-induced hyperemic myocardial blood flow as measured by positron emission tomography: a randomized, double-blind, placebo-controlled crossover trial O. Gaemperli, T. Schepis, P. Koepfli, P. T. Siegrist, S. Fleischman, P. Nguyen, A. Olmsted, W. Wang, H. Lieu and P. A. Kaufmann J Am Coll Cardiol (2008) 51; 328-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18206744
- Application of appropriateness criteria to stress single-photon emission computed tomography sestamibi studies and stress echocardiograms in an academic medical center R. J. Gibbons, T. D. Miller, D. Hodge, L. Urban, P. A. Araoz, P. Pellikka and R. B. McCully J Am Coll Cardiol (2008) 51; 1283-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18371560
- Predictive value of myocardial perfusion single-photon emission computed tomography and the impact of renal function on cardiac death A. Hakeem, S. Bhatti, K. S. Dillie, J. R. Cook, Z. Samad, M. D. Roth-Cline and S. M. Chang Circulation (2008) 118; 2540-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19047585
- NICE, drug-eluting stents and the limits of trial data R. Mohindra Heart (2009) 95; 505-6; author reply 06
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19252009
- Coronary circulatory function abnormalities in insulin resistance: insights from positron emission tomography H. R. Schelbert J Am Coll Cardiol (2009) 53; S3-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19179215
- Interrelation of coronary calcification, myocardial ischemia, and outcomes in patients with intermediate likelihood of coronary artery disease: a combined positron emission tomography/computed tomography study M. P. Schenker, S. Dorbala, E. C. Hong, F. J. Rybicki, R. Hachamovitch, R. Y. Kwong and M. F. Di Carli Circulation (2008) 117;

1693-700

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18362235

A 6 month randomized, double blind, placebo controlled, multi-center trial of high dose atorvastatin on myocardial perfusion abnormalities by positron emission tomography in coronary artery disease S. Sdringola, K. L. Gould, L. G. Zamarka, R. McLain and J. Garner Am Heart J (2008) 155; 245-53

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18215593

Comparison of Troponin T to creatine kinase and to radionuclide cardiac imaging infarct size in patients with ST-elevation myocardial infarction undergoing primary angioplasty D. Tzivoni, D. Koukoui, V. Guetta, L. Novack and G. Cowing Am J Cardiol (2008) 101; 753-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18328834

Prognostic value of multislice computed tomography and gated single-photon emission computed tomography in patients with suspected coronary artery disease J. M. van Werkhoven, J. D. Schuijf, O. Gaemperli, J. W. Jukema, E. Boersma, W. Wijns, P. Stolzmann, H. Alkadhi, I. Valenta, M. P. Stokkel, L. J. Kroft, A. de Roos, G. Pundziute, A. Scholte, E. E. van der Wall, P. A. Kaufmann and J. J. Bax J Am Coll Cardiol (2009) 53; 623-32

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19215839

Role of myocardial perfusion imaging in patients with end-stage renal disease undergoing coronary angiography R. Venkataraman, F. G. Hage, T. Dorfman, J. Heo, R. A. Aqel, A. M. de Mattos and A. E. Iskandrian Am J Cardiol (2008) 102; 1451-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19026294

Comparison of accuracy in the prediction of left ventricular wall motion changes between invasively assessed microvascular integrity indexes and fluorine-18 fluorodeoxyglucose positron emission tomography in patients with ST-elevation myocardial infarction M. H. Yoon, S. J. Tahk, H. M. Yang, S. I. Woo, H. S. Lim, S. J. Kang, B. J. Choi, S. Y. Choi, G. S. Hwang and J. H. Shin Am J Cardiol (2008) 102; 129-34

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18602508