

- Edema as a very early marker for acute myocardial ischemia: a cardiovascular magnetic resonance study H. Abdel-Aty, M. Cocker, C. Meek, J. V. Tyberg and M. G. Friedrich J Am Coll Cardiol (2009) 53; 1194-201
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19341860
- Noninvasive coronary artery imaging: magnetic resonance angiography and multidetector computed tomography angiography: a scientific statement from the american heart association committee on cardiovascular imaging and intervention of the council on cardiovascular radiology and intervention, and the councils on clinical cardiology and cardiovascular disease in the young D. A. Bluemke, S. Achenbach, M. Budoff, T. C. Gerber, B. Gersh, L. D. Hillis, W. G. Hundley, W. J. Manning, B. F. Printz, M. Stuber and P. K. Woodard Circulation (2008) 118; 586-606
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18586979
- Impact of coronary revascularization and transmural extent of scar on regional left ventricular remodelling J. Chan, F. Khafagi, A. A. Young, B. R. Cowan, C. Thompson and T. H. Marwick Eur Heart J (2008) 29; 1608-17
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18556718
- Electroanatomic characterization of post-infarct scars comparison with 3-dimensional myocardial scar reconstruction based on magnetic resonance imaging A. Codreanu, F. Odille, E. Aliot, P. Y. Marie, I. Magnin-Poull, M. Andronache, D. Mandry, W. Djaballah, D. Regent, J. Felblinger and C. de Chillou J Am Coll Cardiol (2008) 52; 839-42
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18755347
- Cardiac magnetic resonance with T2-weighted imaging improves detection of patients with acute coronary syndrome in the emergency department R. C. Cury, K. Shash, J. T. Nagurney, G. Rosito, M. D. Shapiro, C. H. Nomura, S. Abbara, F. Bamberg, M. Ferencik, E. J. Schmidt, D. F. Brown, U. Hoffmann and T. J. Brady Circulation (2008) 118; 837-44
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18678772
- Cardiac magnetic resonance and 64-slice cardiac CT of lipomatous metaplasia of chronic myocardial infarction J. F. Deux, A. Rahmouni and J. Garot Eur Heart J (2008) 29; 570
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17881345
- Techniques to enhance guide catheter support C. Di Mario and N. Ramasami Catheter Cardiovasc Interv (2008) 72; 505-12
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18814225
- The salvaged area at risk in reperfused acute myocardial infarction as visualized by cardiovascular magnetic resonance M. G. Friedrich, H. Abdel-Aty, A. Taylor, J. Schulz-Menger, D. Messroghli and R. Dietz J Am Coll Cardiol (2008) 51; 1581-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18420102
- Predominant subepicardial enhancement on magnetic resonance imaging

corresponding to a post-infarction acute pericarditis in an athletic young patient G. Gahide, R. Gervasoni and F. Roubille Eur Heart J (2008) 29; 2876
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18562302

Cardiac magnetic resonance imaging study for quantification of infarct size comparing directly serial versus single time-point measurements of cardiac troponin T E. Giannitsis, H. Steen, K. Kurz, B. Ivandic, A. C. Simon, S. Futterer, C. Schild, P. Isfort, A. S. Jaffe and H. A. Katus J Am Coll Cardiol (2008) 51; 307-14
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18206741

Relation of left ventricular infarct transmurality and infarct size after primary percutaneous coronary angioplasty to time from symptom onset to balloon inflation J. Y. Hahn, Y. B. Song, H. C. Gwon, Y. H. Choe, J. H. Kim, J. Sung, S. H. Choi, J. H. Choi, D. K. Kim, K. P. Hong, J. E. Park and S. H. Lee Am J Cardiol (2008) 102; 1163-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18940285

Relation between the assessment of microvascular injury by cardiovascular magnetic resonance and coronary Doppler flow velocity measurements in patients with acute anterior wall myocardial infarction A. Hirsch, R. Nijveldt, J. D. Haeck, A. M. Beek, K. T. Koch, J. P. Henriques, R. J. van der Schaaf, M. M. Vis, J. Baan, Jr., R. J. de Winter, J. G. Tijssen, A. C. van Rossum and J. J. Piek J Am Coll Cardiol (2008) 51; 2230-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18534269

Performance of delayed-enhancement magnetic resonance imaging with gadoversetamide contrast for the detection and assessment of myocardial infarction: an international, multicenter, double-blinded, randomized trial R. J. Kim, T. S. Albert, J. H. Wible, M. D. Elliott, J. C. Allen, J. C. Lee, M. Parker, A. Napoli and R. M. Judd Circulation (2008) 117; 629-37
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18212288

Incidence and prognostic implication of unrecognized myocardial scar characterized by cardiac magnetic resonance in diabetic patients without clinical evidence of myocardial infarction R. Y. Kwong, H. Sattar, H. Wu, G. Vorobiof, V. Gandla, K. Steel, S. Siu and K. A. Brown Circulation (2008) 118; 1011-20
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18725488

Relation between stress-induced myocardial perfusion defects on cardiovascular magnetic resonance and coronary microvascular dysfunction in patients with cardiac syndrome X G. A. Lanza, A. Buffon, A. Sestito, L. Natale, G. A. Sgueglia, L. Galiuto, F. Infusino, L. Mariani, A. Centola and F. Crea J Am Coll Cardiol (2008) 51; 466-72
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18222358

Effect of thrombus aspiration on infarct size and left ventricular function in high-risk patients with acute myocardial infarction treated by percutaneous coronary intervention. Results of a prospective controlled pilot study J. Lipiecki, S. Monzy, N. Durel, F. Cachin, P. Chabrot, A. Muliez, D. Morand, J. Maublant and J. Ponsonnaille Am Heart J (2009)

157; 583 e1-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19249433

Use of cardiovascular magnetic resonance imaging in acute coronary syndromes T.

Lockie, E. Nagel, S. Redwood and S. Plein Circulation (2009) 119; 1671-81

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19332480

NICE, drug-eluting stents and the limits of trial data R. Mohindra Heart (2009) 95; 505-6; author reply 06

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19252009

Incremental value of cardiac magnetic resonance imaging in the differential diagnosis of acute coronary syndrome in a young man M. Neizel, M. Kelm and H. P. Kuehl Eur Heart J (2009) 30; 380

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18791201

Functional recovery after acute myocardial infarction: comparison between angiography, electrocardiography, and cardiovascular magnetic resonance measures of microvascular injury R. Nijveldt, A. M. Beek, A. Hirsch, M. G. Stoel, M. B. Hofman, V. A. Umans, P. R. Algra, J. W. Twisk and A. C. van Rossum J Am Coll Cardiol (2008) 52; 181-9

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18617066

A randomized trial of on-pump beating heart and conventional cardioplegic arrest in coronary artery bypass surgery patients with impaired left ventricular function using cardiac magnetic resonance imaging and biochemical markers T. J. Pegg, J. B.

Selvanayagam, J. M. Francis, T. D. Karamitsos, Z. Maunsell, L. M. Yu, S. Neubauer and D. P. Taggart Circulation (2008) 118; 2130-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18981306

Prognostic value of normal adenosine-stress cardiac magnetic resonance imaging G. Pilz, A. Jeske, M. Klos, E. Ali, B. Hoefling, R. Scheck and P. Bernhardt Am J Cardiol (2008) 101; 1408-12

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18471450

Head to head comparison between perfusion and function during accelerated high-dose dipyridamole magnetic resonance stress for the detection of coronary artery disease A. Pingitore, M. Lombardi, B. Scattini, D. De Marchi, G. D. Aquaro, V. Positano and E.

Picano Am J Cardiol (2008) 101; 8-14

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18157957

ST-segment elevation myocardial infarction with concomitant multiple coronary arteries thromboses in a young patient with hyperhomocysteinaemia L. Politi, D. E. Monopoli and M. G. Modena Heart (2008) 94; 1180

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18703695

Usefulness of cardiac magnetic resonance imaging combined with low-dose dobutamine stress to detect an abnormal ventricular stress response in children and young adults after fontan operation at young age D. Robbers-Visser, D. Jan Ten Harkel, L. Kapusta, J. L. Strengers, M. Dalinghaus, F. J. Meijboom, P. M. Pattynama, A. J. Bogers and W. A. Helbing Am J Cardiol (2008) 101; 1657-62
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18489947

Noninvasive detection of fibrosis applying contrast-enhanced cardiac magnetic resonance in different forms of left ventricular hypertrophy relation to remodeling A. Rudolph, H. Abdel-Aty, S. Bohl, P. Boye, A. Zagrosek, R. Dietz and J. Schulz-Menger J Am Coll Cardiol (2009) 53; 284-91
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19147047

Matrix-array 3-dimensional echocardiographic assessment of volumes, mass, and ejection fraction in young pediatric patients with a functional single ventricle: a comparison study with cardiac magnetic resonance B. D. Soriano, M. Hoch, A. Ithurralde, T. Geva, A. J. Powell, B. D. Kussman, D. A. Graham, W. Tworetzky and G. R. Marx Circulation (2008) 117; 1842-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18362236

Hot flashes and subclinical cardiovascular disease: findings from the Study of Women's Health Across the Nation Heart Study R. C. Thurston, K. Sutton-Tyrrell, S. A. Everson-Rose, R. Hess and K. A. Matthews Circulation (2008) 118; 1234-40
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18765392

Effect of rosuvastatin therapy on carotid plaque morphology and composition in moderately hypercholesterolemic patients: a high-resolution magnetic resonance imaging trial H. R. Underhill, C. Yuan, X. Q. Zhao, L. W. Kraiss, D. L. Parker, T. Saam, B. Chu, N. Takaya, F. Liu, N. L. Polissar, B. Neradilek, J. S. Raichlen, V. A. Cain, J. C. Waterton, W. Hamar and T. S. Hatsukami Am Heart J (2008) 155; 584 e1-8
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18294500

Infarct size by contrast enhanced cardiac magnetic resonance is a stronger predictor of outcomes than left ventricular ejection fraction or end-systolic volume index: prospective cohort study E. Wu, J. T. Ortiz, P. Tejedor, D. C. Lee, C. Bucciarelli-Ducci, P. Kansal, J. C. Carr, T. A. Holly, D. Lloyd-Jones, F. J. Klocke and R. O. Bonow Heart (2008) 94; 730-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18070953