

Assessment of coronary flow reserve by coronary pressure measurement: comparison with flow- or velocity-derived coronary flow reserve. Akasaka, T., A. Yamamuro, et al. J Am Coll Cardiol (2003).**41**(9): 1554-60
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12742297

Predictors and implications of residual plaque burden after coronary stenting: an intravascular ultrasound study. Alfonso, F., P. Garcia, et al. Am Heart J (2003).**145**(2): 254-61
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12595842

Myocardial viability, coronary flow reserve, and in-hospital predictors of late recovery of contractility following successful primary stenting for acute myocardial infarction. Beygui, F., C. Le Feuvre, et al. Heart (2003).**89**(2): 179-83
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12527673

Influence of contractile reserve and inducible ischaemia on left ventricular remodelling after acute myocardial infarction. Coletta, C., A. Sestili, et al. Heart (2003).**89**(10): 1138-43
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12975399

Intracoronary and intravenous adenosine 5'-triphosphate, adenosine, papaverine, and contrast medium to assess fractional flow reserve in humans. De Bruyne, B., N. H. Pijls, et al. Circulation (2003).**107**(14): 1877-83
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12668522

Comparison of coronary thermodilution and Doppler velocity for assessing coronary flow reserve. Fearon, W. F., H. M. Farouque, et al. Circulation (2003).**108**(18): 2198-200
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14568891

Novel index for invasively assessing the coronary microcirculation. Fearon, W. F., L. B. Balsam, et al. Circulation (2003).**107**(25): 3129-32
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12821539

Cost-effectiveness of measuring fractional flow reserve to guide coronary interventions. Fearon, W. F., A. C. Yeung, et al. Am Heart J (2003).**145**(5): 882-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12766748

Effect of a change in gender on coronary arterial size: a longitudinal intravascular ultrasound study in transplanted hearts. Herity, N. A., S. Lo, et al. J Am Coll Cardiol (2003).**41**(9): 1539-46
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14568891

[tation&list_uids=12742295](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12742295)

Intravascular ultrasound assessment of patterns of arterial remodeling in the absence of significant reference segment plaque burden in patients with coronary artery disease. Hong, M. K., G. S. Mintz, et al. J Am Coll Cardiol (2003).**42**(5): 806-10

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12957424

Coronary flow velocity analysis during short term follow up after coronary reperfusion: use of transthoracic Doppler echocardiography to predict regional wall motion recovery in patients with acute myocardial infarction.

Hozumi, T., Y. Kanzaki, et al. Heart (2003).**89**(10): 1163-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12975408

Use of fractional flow reserve versus stress perfusion scintigraphy after unstable angina. Effect on duration of hospitalization, cost, procedural characteristics, and clinical outcome. Leesar, M. A., T. Abdul-Baki, et al. J Am Coll Cardiol (2003).**41**(7): 1115-21

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12679210

Measuring pressure-derived fractional flow reserve through four French diagnostic catheters. Legalery, P., M. F. Seronde, et al. Am J Cardiol

(2003).**91**(9): 1075-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12714149

Comparison of transthoracic Doppler echocardiography to intracoronary Doppler guidewire measurements for assessment of coronary flow reserve in the left anterior descending artery for detection of restenosis after coronary angioplasty. Lethen, H., H. P. Tries, et al. Am J Cardiol (2003).**91**(4): 412-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12586254

Impairment of myocardial perfusion in both culprit and nonculprit arteries in acute myocardial infarction: a LIMIT AMI substudy. Murphy, S. A., C. Chen, et al. Am J Cardiol (2003).**91**(3): 325-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12565089

Role of incremental doses of intracoronary adenosine for fractional flow reserve assessment. Murtagh, B., S. Higano, et al. Am Heart J (2003).**146**(1): 99-105

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12851614

Persistence of systolic coronary flow reversal predicts irreversible dysfunction after reperfused anterior myocardial infarction. Nohtomi, Y., M. Takeuchi, et al. Heart (2003).**89**(4): 382-8

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Ci>

[tation&list_uids=12639863](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12639863)

Assessment of myocardial viability using coronary zero flow pressure after successful angioplasty in patients with acute anterior myocardial infarction.

Shimada, K., Y. Sakanoue, et al. *Heart* (2003).**89**(1): 71-6

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12482796

Phasic coronary blood flow velocity pattern and flow reserve in the atrium: regulation of left atrial myocardial perfusion. Skalidis, E. I., G. E. Kochiadakis, et al. *J Am Coll Cardiol* (2003).**41**(4): 674-80

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12598082

Reliability of pressure-derived myocardial fractional flow reserve in assessing coronary artery stenosis in patients with previous myocardial infarction. Usui, Y., T. Chikamori, et al. *Am J Cardiol* (2003).**92**(6): 699-702

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12972110