

**Clinical and angiographic outcome of directional atherectomy followed by stent implantation in de novo lesions located at the ostium of the left anterior descending coronary artery.** Airoidi, F., C. Di Mario, et al. Heart (2003).**89**(9): 1050-4

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12923025](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12923025)

**Impact of intracoronary radiation on in-stent restenosis involving ostial lesions.** Ajani, A. E., R. Waksman, et al. Catheter Cardiovasc Interv (2003).**58**(2): 175-80

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12552539](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12552539)

**Intravascular brachytherapy for native coronary ostial in-stent restenotic lesions.** Costantini, C. O., A. J. Lansky, et al. J Am Coll Cardiol (2003).**41**(10): 1725-31

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12767655](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12767655)