

Blockade of the intermediate-conductance calcium-activated potassium channel as a new therapeutic strategy for restenosis. Kohler, R., H. Wulff, et al. Circulation (2003).**108**(9): 1119-25

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12939222

Impact of deep vessel wall injury on acute response and remodeling of coronary artery segments after cutting balloon angioplasty. Nakamura, M., P. G. Yock, et al. Am J Cardiol (2003).**91**(1): 6-11

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12505563

Single injection of P-selectin or P-selectin glycoprotein ligand-1 monoclonal antibody blocks neointima formation after arterial injury in apolipoprotein E-deficient mice. Phillips, J. W., K. G. Barrinhaus, et al. Circulation (2003).**107**(17): 2244-9

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12707243

Elevated whole-blood tissue factor procoagulant activity as a marker of restenosis after percutaneous transluminal coronary angioplasty and stent implantation. Tutar, E., M. Ozcan, et al. Circulation (2003).**108**(13): 1581-4

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12975255

Clinical outcomes of stents versus balloon angioplasty in non-acute coronary artery disease. A meta-analysis of randomized controlled trials.

Nordmann, A. J., P. Hengstler, et al. Eur Heart J (2004).**25**(1): 69-80

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14683745