

**Effect of recanalization of chronic total occlusions on global and regional left ventricular function in patients with or without previous myocardial infarction.** Chung, C. M., S. Nakamura, et al. Catheter Cardiovasc Interv (2003).**60**(3): 368-74

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=14571489](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14571489)

**Intracoronary beta-brachytherapy in chronic total occlusions: a subgroup analysis from the RENO registry.** Jain, D., V. Geist, et al. Catheter Cardiovasc Interv (2003).**58**(3): 322-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12594695](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12594695)

**Immediate results and one-year clinical outcome after percutaneous coronary interventions in chronic total occlusions: data from a multicenter, prospective, observational study (TOAST-GISE).** Olivari, Z., P. Rubartelli, et al. J Am Coll Cardiol (2003).**41**(10): 1672-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12767645](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12767645)

**Coronary stent implantation is superior to balloon angioplasty for chronic coronary occlusions: six-year clinical follow-up of the GISSOC trial.** Rubartelli, P., E. Verna, et al. J Am Coll Cardiol (2003).**41**(9): 1488-92

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12742287](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12742287)

**Angioplasty for chronic total occlusion by using tapered-tip guidewires.** Saito, S., S. Tanaka, et al. Catheter Cardiovasc Interv (2003).**59**(3): 305-11

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12822146](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12822146)

**Incidence and predictors of late total occlusion following coronary stenting.**

Shah, P. B., D. E. Cutlip, et al. Catheter Cardiovasc Interv (2003).**60**(3): 344-51  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=14571485](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14571485)

**Angiographic assessment of collateral connections in comparison with invasively determined collateral function in chronic coronary occlusions.**

Werner, G. S., M. Ferrari, et al. Circulation (2003).**107**(15): 1972-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12665484](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12665484)

**Determinants of target vessel failure in chronic total coronary occlusions after stent implantation. The influence of collateral function and coronary hemodynamics.** Werner, G. S., P. Bahrmann, et al. J Am Coll Cardiol (2003).**42**(2): 219-25

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=12875755](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12875755)