

Survival and cardiac remodeling benefits in patients undergoing late percutaneous coronary intervention of the infarct-related artery: evidence from a meta-analysis of randomized controlled trials A. Abbate, G. G. Biondi-Zoccai, D. L. Appleton, P. Erne, A. W. Schoenenberger, M. J. Lipinski, P. Agostoni, I. Sheiban and G. W. Vetrovec *J Am Coll Cardiol* (2008) 51; 956-64

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18308165](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18308165)

Percutaneous intervention for chronic total occlusion of the internal iliac artery for unrelenting buttock claudication S. Adlakhia, M. Burket and C. Cooper *Catheter Cardiovasc Interv* (2008)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19213070](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19213070)

Culotte stenting technique in coronary bifurcation disease: angiographic follow-up using dedicated quantitative coronary angiographic analysis and 12-month clinical outcomes T. Adriaenssens, R. A. Byrne, A. Dibra, R. Iijima, J. Mehilli, O. Bruskin, A. Schomig and A. Kastrati *Eur Heart J* (2008) 29; 2868-76

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19001472](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19001472)

Rural interhospital transfer of ST-elevation myocardial infarction patients for percutaneous coronary revascularization: the Stat Heart Program F. V. Aguirre, J. J. Varghese, M. P. Kelley, W. Lam, C. L. Lucre, J. B. Gill, L. Page, L. Turner, C. Davis and F. L. Mikell *Circulation* (2008) 117; 1145-52

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18268151](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18268151)

Treatment of acute limb ischemia with a percutaneous mechanical thrombectomy-based endovascular approach: 5-year limb salvage and survival results from a single center series G. M. Ansel, C. F. Botti, Jr. and M. J. Silver *Catheter Cardiovasc Interv* (2008) 72; 325-30

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18726955](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18726955)

Early and late benefits of prasugrel in patients with acute coronary syndromes undergoing percutaneous coronary intervention: a TRITON-TIMI 38 (TRial to Assess Improvement in Therapeutic Outcomes by Optimizing Platelet Inhibition with Prasugrel-Thrombolysis In Myocardial Infarction) analysis E. M. Antman, S. D. Wiviott, S. A. Murphy, J. Voitek, Y. Hasin, P. Widimsky, H. Chandna, W. Macias, C. H. McCabe and E. Braunwald *J Am Coll Cardiol* (2008) 51; 2028-33

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18498956](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18498956)

Late percutaneous coronary intervention for the totally occluded infarct-related artery: a meta-analysis of the effects on cardiac function and remodeling D. L. Appleton, A. Abbate and G. G. Biondi-Zoccai *Catheter Cardiovasc Interv* (2008) 71; 772-81

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18415952](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18415952)

Feasibility of primary clot extraction prior to percutaneous coronary intervention in acute myocardial infarction R. Aqel, G. Zoghbi, F. Hage, G. Philips, G. Perry, A. Iskandrian and L. Dell'Italia *Catheter Cardiovasc Interv* (2008) 71; 870-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18383164](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18383164)

Medical therapy versus percutaneous coronary intervention for patients with stable coronary artery disease A. Arbab-Zadeh *J Am Coll Cardiol* (2009) 53; 528-9; author reply 29

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19195616](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195616)

Effect of intravenous FX06 as an adjunct to primary percutaneous coronary intervention for acute ST-segment elevation myocardial infarction results of the F.I.R.E. (Efficacy of FX06 in the Prevention of Myocardial Reperfusion Injury) trial D. Atar, P. Petzelbauer, J. Schwitter, K. Huber, B. Rensing, J. D. Kasprzak, C. Butter, L. Grip, P. R. Hansen, T. Suselbeck, P. M. Clemmensen, M. Marin-Galiano, B. Geudelin and P. T. Buser *J Am Coll Cardiol* (2009) 53; 720-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19232907](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19232907)

Predictors of slow flow during primary percutaneous coronary intervention: an intravascular ultrasound-virtual histology study J. H. Bae, T. G. Kwon, D. W. Hyun, C. S. Rihal and A. Lerman *Heart* (2008) 94; 1559-64

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18381376](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18381376)

Percutaneous intervention on the saphenous vein bypass grafts--long-term outcomes D. Bansal, R. Muppidi, S. Singla, R. Sukhija, S. Zarich, J. L. Mehta and R. Sachdeva *Catheter Cardiovasc Interv* (2008) 71; 58-61

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18098183](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18098183)

Percutaneous intervention in saphenous vein bypass graft disease: case against the use of drug-eluting stents D. Bansal, R. Sachdeva and J. L. Mehta *J Am Coll Cardiol* (2008) 51; 970-1; author reply 71-2

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18308169](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18308169)

Optical coherence tomography assessment of vulnerable plaque rupture: predilection for the plaque 'shoulder' P. Barlis, P. W. Serruys, A. Devries and E. Regar *Eur Heart J* (2008) 29; 2023

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18337236](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18337236)

Intracoronary KAI-9803 as an adjunct to primary percutaneous coronary intervention for acute ST-segment elevation myocardial infarction E. Bates, C. Bode, M. Costa, C. M. Gibson, C. Granger, C. Green, K. Grimes, R. Harrington, K. Huber, N. Kleiman, D. Mochly-Rosen, M. Roe, Z. Sadowski, S. Solomon and P. Widimsky *Circulation* (2008) 117; 886-96

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18250271](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18250271)

Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a comprehensive meta-analysis of randomized trials A. A. Bavry, D. J. Kumbhani and D. L. Bhatt *Eur Heart J* (2008) 29; 2989-3001

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list\_uids=18812323

Survival benefit after percutaneous treatment of chronic total coronary occlusions G. Belli *Eur Heart J* (2009) 30; 506; author reply 06-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19168525](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19168525)

Percutaneous coronary intervention in nonagenarians: age before beauty? D. L. Bhatt *Catheter Cardiovasc Interv* (2008) 71; 531-2

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18307234](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18307234)

Emergence of the Concept of Platelet Reactivity Monitoring of Response to Thienopyridines L. Bonello, A. De Labriolle, M. Scheinowitz, G. Lemesle, P. Roy, D. H. Steinberg, T. L. Pinto Slottow, R. Pakala, A. D. Pichard, P. Barragan, L. Camoin-Jau, F. Dignat-George, F. Paganelli and R. Waksman *Heart* (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19196732](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19196732)

Thrombus aspiration during percutaneous coronary intervention L. Bonello, A. De Labriolle, D. Steinberg and R. Waksman *Lancet* (2008) 372; 1034; author reply 34-5

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18805323](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18805323)

Randomized, controlled trial of coronary artery bypass surgery versus percutaneous coronary intervention in patients with multivessel coronary artery disease: six-year follow-up from the Stent or Surgery Trial (SoS) J. Booth, T. Clayton, J. Pepper, F. Nugara, M. Flather, U. Sigwart and R. H. Stables *Circulation* (2008) 118; 381-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18606919](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18606919)

Treatment of coronary bifurcations: might less be more? E. S. Brilakis and P. B. Berger *Eur Heart J* (2008) 29; 704-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18343799](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18343799)

Transient and persistent renal dysfunction are predictors of survival after percutaneous coronary intervention: insights from the Dartmouth Dynamic Registry J. R. Brown, D. J. Malenka, J. T. DeVries, J. F. Robb, J. E. Jayne, B. J. Friedman, B. D. Hettleman, N. W. Niles, A. V. Kaplan, A. C. Schoolwerth and C. A. Thompson *Catheter Cardiovasc Interv* (2008) 72; 347-54

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18729173](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18729173)

Do patients with angina alone have a more benign prognosis than patients with a history of acute myocardial infarction, revascularisation or both? Findings from a community cohort study B. Buckley and A. W. Murphy *Heart* (2009) 95; 461-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18669551](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18669551)

ST-segment recovery and outcome after primary percutaneous coronary intervention for ST-elevation myocardial infarction: insights from the Assessment of Pexelizumab in Acute Myocardial Infarction (APEX-AMI) trial C. E. Buller, Y. Fu, K. W. Mahaffey, T. G. Todaro, P. Adams, C. M. Westerhout, H. D. White, A. W. van 't Hof, F. J. Van de Werf, G. S. Wagner, C. B. Granger and P. W. Armstrong *Circulation* (2008) 118; 1335-46

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18779444](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18779444)

Percutaneous versus surgical revascularization for multivessel coronary artery disease: A single center 10 year follow-up of SOS trial patients P. Buszman, S. Wiernek, R. Szymanski, B. Bialkowska, W. Fil, R. Stables, A. Bochenek, J. Martin and M. Tendera *Catheter Cardiovasc Interv* (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19360866](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19360866)

Coronary stent fracture: the hidden truth of a problem more common than stent thrombosis J. D. Carroll *Catheter Cardiovasc Interv* (2009) 73; 88-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19089948](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19089948)

Time-to-reperfusion in patients undergoing interhospital transfer for primary percutaneous coronary intervention in the U.S: an analysis of 2005 and 2006 data from the National Cardiovascular Data Registry A. Chakrabarti, H. M. Krumholz, Y. Wang, J. S. Rumsfeld and B. K. Nallamothu *J Am Coll Cardiol* (2008) 51; 2442-3

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18565404](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18565404)

The revascular active percutaneous interventional device for coronary total occlusions study D. Chamie, A. Abizaid, J. R. Costa, Jr., F. Feres, R. Staico, R. Costa, L. A. Mattos, A. G. Sousa and J. E. Sousa *Catheter Cardiovasc Interv* (2008) 72; 156-63

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18655121](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18655121)

Predictors of initial nontherapeutic anticoagulation with unfractionated heparin in ST-segment elevation myocardial infarction S. Cheng, D. A. Morrow, S. Sloan, E. M. Antman and M. S. Sabatine *Circulation* (2009) 119; 1195-202

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19237657](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19237657)

Percutaneous treatment of a popliteal bifurcation--the value of coronary devices and strategies R. Colantonio, A. Latib and G. M. Sangiorgi *Catheter Cardiovasc Interv* (2008) 72; 710-3

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18949797](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18949797)

Cytochrome P450 2C19 polymorphism in young patients treated with clopidogrel after myocardial infarction: a cohort study J. P. Collet, J. S. Hulot, A. Pena, E. Villard, J. B. Esteve, J. Silvain, L. Payot, D. Brugier, G. Cayla, F. Beygui, G. Bensimon, C. Funck-Brentano and G. Montalescot *Lancet* (2009) 373; 309-17

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19108880](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19108880)

Long-term safety and efficacy of percutaneous coronary intervention with stenting and coronary artery bypass surgery for multivessel coronary artery disease: a meta-analysis with 5-year patient-level data from the ARTS, ERACI-II, MASS-II, and SoS trials J. Daemen, E. Boersma, M. Flather, J. Booth, R. Stables, A. Rodriguez, G. Rodriguez-Granillo, W. A. Hueb, P. A. Lemos and P. W. Serruys *Circulation* (2008) 118; 1146-54

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list\_uids=18725490

PCI at non-PCI centres: immediate or rescue? N. Danchin and P. W. Armstrong *Lancet* (2008) 371; 534-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18280311](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18280311)

Comparison of thrombolysis followed by broad use of percutaneous coronary intervention with primary percutaneous coronary intervention for ST-segment-elevation acute myocardial infarction: data from the french registry on acute ST-elevation myocardial infarction (FAST-MI) N. Danchin, P. Coste, J. Ferrieres, P. G. Steg, Y. Cottin, D. Blanchard, L. Belle, B. Ritz, G. Kirkorian, M. Angioi, P. Sans, B. Charbonnier, H. Eltchaninoff, P. Gueret, K. Khalife, P. Asseman, J. Puel, P. Goldstein, J. P. Cambou and T. Simon *Circulation* (2008) 118; 268-76

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18591434](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18591434)

Evaluating risks and benefits in coronary revascularisation--a very imperfect art? M. A. de Belder and L. Hamilton *Heart* (2009) 95; 6-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18768566](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18768566)

Early glycoprotein IIb-IIIa inhibitors in primary angioplasty (EGYPT) cooperation: an individual patient data meta-analysis G. De Luca, C. M. Gibson, F. Bellandi, S. Murphy, M. Maioli, M. Noc, U. Zeymer, D. Dudek, H. R. Arntz, S. Zorman, H. M. Gabriel, A. Emre, D. Cutlip, G. Biondi-Zoccai, T. Rakowski, M. Gyongyosi, P. Marino, K. Huber and A. W. van't Hof *Heart* (2008) 94; 1548-58

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18474534](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18474534)

Vascular calcification: pathobiology of a multifaceted disease L. L. Demer and Y. Tintut *Circulation* (2008) 117; 2938-48

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18519861](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18519861)

Anatomic distribution of the culprit lesion in patients with non-ST-segment elevation myocardial infarction undergoing percutaneous coronary intervention: findings from the National Cardiovascular Data Registry W. C. t. Dixon, T. Y. Wang, D. Dai, K. A. Shunk, E. D. Peterson and M. T. Roe *J Am Coll Cardiol* (2008) 52; 1347-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18929247](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18929247)

Percutaneous coronary intervention for chronic total occlusion: novel 3-dimensional imaging and quantitative analysis D. Dvir, A. Assali and R. Kornowski *Catheter Cardiovasc Interv* (2008) 71; 784-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18415954](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18415954)

Trends in acute reperfusion therapy for ST-segment elevation myocardial infarction from 1999 to 2006: we are getting better but we have got a long way to go K. A. Eagle, B. K. Nallamothu, R. H. Mehta, C. B. Granger, P. G. Steg, F. Van de Werf, J. Lopez-Sendon, S. G. Goodman, A. Quill and K. A. Fox *Eur Heart J* (2008) 29; 609-17

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18310671](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18310671)

Refining the art and science of coronary stenting S. G. Ellis *N Engl J Med* (2009) 360; 292-4

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19144945](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19144945)

Safety and efficacy of drug-eluting balloons in percutaneous treatment of bifurcation lesions: the DEBIUT (drug-eluting balloon in bifurcation Utrecht) registry J. C.

Fanggiday, P. R. Stella, S. H. Guyomi and P. A. Doevendans *Catheter Cardiovasc Interv* (2008) 71; 629-35

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18360855](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18360855)

Incidence, predictors, and outcome of reinfarction and stent thrombosis within one year after primary percutaneous coronary intervention for ST-elevation myocardial infarction

M. L. Fokkema, P. A. van der Vleuten, P. J. Vlaar, T. Svilaas and F. Zijlstra *Catheter Cardiovasc Interv* (2009) 73; 627-34

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19309712](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19309712)

Incidence and clinical consequences of distal embolization on the coronary angiogram after percutaneous coronary intervention for ST-elevation myocardial infarction

M. L. Fokkema, P. J. Vlaar, T. Svilaas, M. Vogelzang, D. Amo, G. F. Diercks, A. J. Suurmeijer and F. Zijlstra *Eur Heart J* (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19224928](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19224928)

Outcomes after percutaneous coronary intervention of ostial lesions in the era of drug-eluting stents

M. Freeman, D. J. Clark, N. Andrianopoulos, S. J. Duffy, H. S. Lim, A. Brennan, K. Charter, J. Shaw, M. Horrigan, A. E. Ajani, M. Sebastian, C. M. Reid and H. M. Farouque *Catheter Cardiovasc Interv* (2008)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19309731](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19309731)

Abbreviated infusion of eptifibatide after successful coronary intervention The BRIEF-PCI (Brief Infusion of Eptifibatide Following Percutaneous Coronary Intervention)

randomized trial A. Y. Fung, J. Saw, A. Starovoytov, C. Densem, P. Jokhi, S. J. Walsh, R. S. Fox, K. H. Humphries, E. Aymong, D. R. Ricci, J. G. Webb, J. N. Hamburger, R. G. Carere and C. E. Buller *J Am Coll Cardiol* (2009) 53; 837-45

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19264239](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19264239)

Impact of anticoagulation regimens on sheath management and bleeding in patients undergoing elective percutaneous coronary intervention in the STEEPLE trial

R. Gallo, S. R. Steinhubl, H. D. White and G. Montalescot *Catheter Cardiovasc Interv* (2009) 73; 319-25

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19213086](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19213086)

The effect of percutaneous coronary intervention on inflammatory response and endothelial progenitor cell recruitment

R. Garg, A. Tellez, C. Alviar, J. Granada, N. S. Kleiman and E. I. Lev *Catheter Cardiovasc Interv* (2008) 72; 205-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18651648](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18651648)

Suspected coronary artery dissection post-stenting, confirmed by optical coherence tomography S. Garg, C. Bourantas and S. Thackray *Heart* (2008) 94; 335  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18276819](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18276819)

Acute reperfusion strategies for ST-segment elevation myocardial infarction B. M. Glover and A. A. Adgey *Eur Heart J* (2008) 29; 571-2  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18310670](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18310670)

Clot removal in AMI--worth the hassle? D. A. Gorog and I. Malik *Heart* (2008) 94; 258-9  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18276807](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18276807)

The relative safety and efficacy of abciximab and eptifibatid in patients undergoing primary percutaneous coronary intervention: insights from a large regional registry of contemporary percutaneous coronary intervention H. S. Gurm, D. E. Smith, J. S. Collins, D. Share, A. Riba, A. J. Carter, T. LaLonde, E. Kline-Rogers, M. O'Donnell, H. Changezi, M. Zughuib, R. Safian and M. Moscucci *J Am Coll Cardiol* (2008) 51; 529-35  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18237680](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18237680)

Percutaneous coronary intervention in anticoagulated patients via radial artery access G. Helft, G. Dambrin, A. Zaman, C. Le Feuvre, O. Barthelemy, F. Beygui, X. Favereau and J. P. Metzger *Catheter Cardiovasc Interv* (2009) 73; 44-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19089936](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19089936)

Successful percutaneous closure of coronary artery fistula with angiographic follow-up at 6 months C. Hendry, V. Mahadevan and F. Fath-Ordoubadi *Catheter Cardiovasc Interv* (2009) 73; 581-3  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19085916](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19085916)

Carotid angioplasty and stenting in octogenarians: is it safe? M. Henry, I. Henry, A. Polydorou and M. Hugel *Catheter Cardiovasc Interv* (2008) 72; 309-17  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18729151](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18729151)

Impact of plaque components on no-reflow phenomenon after stent deployment in patients with acute coronary syndrome: a virtual histology-intravascular ultrasound analysis Y. J. Hong, M. H. Jeong, Y. H. Choi, J. S. Ko, M. G. Lee, W. Y. Kang, S. E. Lee, S. H. Kim, K. H. Park, D. S. Sim, N. S. Yoon, H. J. Youn, K. H. Kim, H. W. Park, J. H. Kim, Y. Ahn, J. G. Cho, J. C. Park and J. C. Kang *Eur Heart J* (2009)  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19228713](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19228713)

Traditional antegrade approach versus combined antegrade and retrograde approach in the percutaneous treatment of coronary chronic total occlusions J. T. Hsu, H. Tamai, E. Kyo, T. Tsuji and S. Watanabe *Catheter Cardiovasc Interv* (2009)  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19360876](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19360876)

Cardiovascular events and re-stenosis following administration of G-CSF in acute myocardial infarction: systematic review and meta-analysis H. Ince, M. Valgimigli, M.

Petzsch, J. S. de Lezo, F. Kuethe, S. Dunkelmann, G. Biondi-Zoccai and C. A. Nienaber  
Heart (2008) 94; 610-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17761504](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17761504)

Role of percutaneous coronary minimalist intervention in the management of acute ST-segment elevation myocardial infarction K. Isaz J Am Coll Cardiol (2008) 51; 1046-7; author reply 47

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18325448](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18325448)

Neointimal hyperplasia after sirolimus-eluting and paclitaxel-eluting stent implantation in diabetic patients: the Randomized Diabetes and Drug-Eluting Stent (DiabeDES) Intravascular Ultrasound Trial L. O. Jensen, M. Maeng, P. Thayssen, E. H. Christiansen, K. N. Hansen, A. Galloe, H. Kelbaek, J. F. Lassen and L. Thuesen Eur Heart J (2008) 29; 2733-41

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18832385](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18832385)

Prevalence and prognostic significance of preprocedural cardiac troponin elevation among patients with stable coronary artery disease undergoing percutaneous coronary intervention: results from the evaluation of drug eluting stents and ischemic events registry A. Jeremias, N. S. Kleiman, D. Nassif, W. H. Hsieh, M. Pencina, K. Maresh, M. Parikh, D. E. Cutlip, R. Waksman, S. Goldberg, P. B. Berger and D. J. Cohen Circulation (2008) 118; 632-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18645057](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18645057)

Paclitaxel-eluting stent long-term outcomes in percutaneous saphenous vein graft interventions (PELOPS) study M. H. Jim, H. H. Ho, R. L. Ko, K. H. Yiu, C. W. Siu, C. P. Lau and W. H. Chow Am J Cardiol (2009) 103; 199-202

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19121436](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19121436)

Effects of aspirin dose on ischaemic events and bleeding after percutaneous coronary intervention: insights from the PCI-CURE study S. S. Jolly, J. Pogue, K. Haladyn, R. J. Peters, K. A. Fox, A. Avezum, B. J. Gersh, H. J. Rupprecht, S. Yusuf and S. R. Mehta Eur Heart J (2008)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18819961](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18819961)

Thrombogenicity in patients with percutaneous coronary artery intervention and dual antiplatelet treatment P. W. Kamphuisen Eur Heart J (2008) 29; 1699-700

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18544543](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18544543)

Multicenter randomized trial of facilitated percutaneous coronary intervention with low-dose tenecteplase in patients with acute myocardial infarction: The Athens PCI trial J. Kanakakis, J. N. Nanas, E. P. Tsagalou, G. D. Maroulidis, S. G. Drakos, A. S. Ntalianis, P. Tzoumele, E. Skoumbourdis, P. Charbis, S. Rokas and M. Anastasiou-Nana Catheter Cardiovasc Interv (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19360864](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19360864)



Safety of percutaneous coronary intervention during uninterrupted oral anticoagulant treatment P. P. Karjalainen, S. Vikman, M. Niemela, P. Porela, A. Ylitalo, M. A. Vaittinen, M. Puurunen, T. J. Airaksinen, K. Nyman, T. Vahlberg and K. E. Airaksinen *Eur Heart J* (2008) 29; 1001-10

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18346963](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18346963)

Percutaneous coronary intervention for stable coronary artery disease D. G. Katriasis and B. Meier *J Am Coll Cardiol* (2008) 52; 889-93

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18772057](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18772057)

Intravenous administration of nicorandil immediately before percutaneous coronary intervention can prevent slow coronary flow phenomenon Y. Kawai, K. Hisamatsu, H. Matsubara, K. Dan, S. Akagi, K. Miyaji, M. Munemasa, Y. Fujimoto, K. F. Kusano and T. Ohe *Eur Heart J* (2009) 30; 765-72

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19276198](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19276198)

Randomized comparison of distal protection versus conventional treatment in primary percutaneous coronary intervention: the drug elution and distal protection in ST-elevation myocardial infarction (DEDICATION) trial H. Kelbaek, C. J. Terkelsen, S. Helqvist, J. F. Lassen, P. Clemmensen, L. Klovgaard, A. Kaltoft, T. Engstrom, H. E. Botker, K. Saunamaki, L. R. Krusell, E. Jorgensen, H. H. Hansen, E. H. Christiansen, J. Ravkilde, L. Kober, K. F. Kofoed and L. Thuesen *J Am Coll Cardiol* (2008) 51; 899-905

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18308157](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18308157)

Use of the Stereotaxis Niobe magnetic navigation system for percutaneous coronary intervention: results from 350 consecutive patients F. Kiemeneij, M. S. Patterson, G. Amoroso, G. Laarman and T. Slagboom *Catheter Cardiovasc Interv* (2008) 71; 510-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18307222](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18307222)

Long-term outcomes of coronary-artery bypass graft surgery versus percutaneous coronary intervention for multivessel coronary artery disease in the bare-metal stent era T. Kimura, T. Morimoto, Y. Furukawa, Y. Nakagawa, S. Shizuta, N. Ehara, R. Taniguchi, T. Doi, K. Nishiyama, N. Ozasa, N. Saito, K. Hoshino, H. Mitsuoka, M. Abe, M. Toma, T. Tamura, Y. Haruna, Y. Imai, S. Teramukai, M. Fukushima and T. Kita *Circulation* (2008) 118; S199-209

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18824755](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18824755)

Serum creatinine ratio: A novel predictor of mortality after percutaneous coronary intervention in patients with normal and abnormal renal function A. S. Kini, K. Sarkar, O. C. Rafael, M. Jakkula, D. Kaplish, P. Lee, J. Suleman, P. Krishnan, M. C. Kim and S. K. Sharma *Catheter Cardiovasc Interv* (2008)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19288579](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19288579)

The problem with composite end points in cardiovascular studies: the story of major adverse cardiac events and percutaneous coronary intervention K. E. Kip, K.

Hollabaugh, O. C. Marroquin and D. O. Williams *J Am Coll Cardiol* (2008) 51; 701-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18279733](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18279733)

Safety and tolerability of hirudins in percutaneous coronary intervention and the risk of Kounis syndrome N. G. Kounis, S. A. Kounis, G. N. Kounis and G. Hahalis *Catheter Cardiovasc Interv* (2009) 73; 584-5

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19085914](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19085914)

Presence of older thrombus is an independent predictor of long-term mortality in patients with ST-elevation myocardial infarction treated with thrombus aspiration during primary percutaneous coronary intervention M. C. Kramer, A. C. van der Wal, K. T. Koch, J. P. Ploegmakers, R. J. van der Schaaf, J. P. Henriques, J. Baan, Jr., S. Z. Rittersma, M. M. Vis, J. J. Piek, J. G. Tijssen and R. J. de Winter *Circulation* (2008) 118; 1810-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18852369](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18852369)

External validation of established risk adjustment models for procedural complications after percutaneous coronary intervention B. Kunadian, J. Dunning, R. Das, A. P. Roberts, R. Morley, A. J. Turley, D. Twomey, J. A. Hall, R. A. Wright, A. G. Sutton, D. F. Muir and M. A. de Belder *Heart* (2008) 94; 1012-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18032457](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18032457)

Concordant improvements in coronary flow reserve and ST-segment resolution during percutaneous coronary intervention for acute myocardial infarction: a benefit of postconditioning W. K. Laskey, S. Yoon, N. Calzada and M. J. Ricciardi *Catheter Cardiovasc Interv* (2008) 72; 212-20

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18546233](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18546233)

Two stent or not two stent--that is the question J. F. Lassen *Eur Heart J* (2008) 29; 2829-30

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18988668](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18988668)

Multicenter international registry of unprotected left main coronary artery percutaneous coronary intervention with drug-eluting stents in patients with myocardial infarction M. S. Lee, D. Sillano, A. Latib, A. Chieffo, G. B. Zoccai, R. Bhatia, I. Sheiban, A. Colombo and J. Tobis *Catheter Cardiovasc Interv* (2009) 73; 15-21

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19089930](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19089930)

Outcomes of nonagenarians who undergo percutaneous coronary intervention with drug-eluting stents M. S. Lee, R. Zimmer, A. Pessegueiro, D. Jurewitz and J. Tobis *Catheter Cardiovasc Interv* (2008) 71; 526-30

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18307230](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18307230)

Impact of bivalirudin on in-hospital bleeding and six-month outcomes in octogenarians undergoing percutaneous coronary intervention G. Lemesle, A. De Labriolle, L. Bonello, A. Syed, S. Collins, G. Maluenda, R. Torguson, K. Kaneshige, Z. Xue, W. O. Suddath, L. F. Satler, K. M. Kent, J. Lindsay, A. D. Pichard and R. Waksman *Catheter Cardiovasc*

Interv (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19360860](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19360860)

Left ventricular assist via percutaneous transhepatic transseptal cannulation in swine D. S. Lim, C. J. Cortese, A. N. Loree, D. A. Dean and R. G. Svittek *Catheter Cardiovasc Interv* (2008)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19213076](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19213076)

Percutaneous transthoracic ventricular puncture for diagnostic and interventional catheterization D. S. Lim, M. Ragosta and J. M. Dent *Catheter Cardiovasc Interv* (2008) 71; 915-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18383174](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18383174)

Prognostic value of ST-segment resolution after rescue percutaneous coronary intervention. Data from the RICO survey L. Lorgis, M. Zeller, G. Dentan, Y. Laurent, J. A. Taam, I. L'Huillier, M. Vincent-Martin, H. Makki and Y. Cottin *Catheter Cardiovasc Interv* (2008) 71; 607-12

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18360851](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18360851)

Diagnostic-therapeutic cascade revisited: coronary angiography, coronary artery bypass graft surgery, and percutaneous coronary intervention in the modern era F. L. Lucas, A. E. Siewers, D. J. Malenka and D. E. Wennberg *Circulation* (2008) 118; 2797-802

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19064681](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19064681)

Assessing the risks of percutaneous coronary intervention: do we have an equivalent of the EuroSCORE? P. F. Ludman *Heart* (2008) 94; 1366-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18931150](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18931150)

Anterior myocardial infarction with acute percutaneous coronary intervention and intracoronary injection of autologous mononuclear bone marrow cells: safety, clinical outcome, and serial changes in left ventricular function during 12-months' follow-up K. Lunde, S. Solheim, K. Forfang, H. Arnesen, L. Brinch, R. Bjornerheim, A. Ragnarsson, T. Egeland, K. Endresen, A. Ilebekk, A. Mangschau and S. Aakhus *J Am Coll Cardiol* (2008) 51; 674-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18261689](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18261689)

Clinical impact of an inter-hospital transfer strategy in patients with ST-elevation myocardial infarction undergoing primary angioplasty: the Emilia-Romagna ST-segment elevation acute myocardial infarction network A. Manari, P. Ortolani, P. Guastaroba, G. Casella, L. Vignali, E. Varani, G. Piovaccari, V. Guiducci, G. Percoco, S. Tondi, F. Passerini, A. Santarelli and A. Marzocchi *Eur Heart J* (2008) 29; 1834-42

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18617475](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18617475)

Percutaneous removal of embolized port catheter: The standard option P. Y. Marcy, J. Thariat and N. Amoretti *Catheter Cardiovasc Interv* (2008) 72; 1025-6

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list\_uids=19021275

Quality of life after late invasive therapy for occluded arteries D. B. Mark, W. Pan, N. E. Clapp-Channing, K. J. Anstrom, J. R. Ross, R. S. Fox, G. P. Devlin, C. E. Martin, C. Adlbrecht, P. A. Cowper, L. D. Ray, E. A. Cohen, G. A. Lamas and J. S. Hochman *N Engl J Med* (2009) 360; 774-83

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19228620](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19228620)

Long-term mortality after bolus-only administration of abciximab, eptifibatide, or tirofiban during percutaneous coronary intervention J. D. Marmur, S. Poludasu, J. Lazar and E. Cavusoglu *Catheter Cardiovasc Interv* (2009) 73; 214-21

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19156882](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19156882)

A randomized trial comparing compression, Perclose Proglide and Angio-Seal VIP for arterial closure following percutaneous coronary intervention: the CAP trial J. L. Martin, A. Pratsos, E. Magargee, K. Mayhew, C. Pensyl, M. Nunn, F. Day and T. Shapiro *Catheter Cardiovasc Interv* (2008) 71; 1-5

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18098171](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18098171)

Failed percutaneous coronary intervention: a decade of experience in 21,000 patients S. J. Mattichak, S. R. Dixon, F. Shannon, J. A. Boura and R. D. Safian *Catheter Cardiovasc Interv* (2008) 71; 131-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18231989](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18231989)

Determinants of coronary arterial flow-mediated dilatation following percutaneous coronary intervention M. McGrady, P. Thanyasiri, B. P. Bailey, D. S. Celermajer and M. R. Adams *Catheter Cardiovasc Interv* (2008) 71; 327-32

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18288727](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18288727)

Abciximab in Patients With Acute ST-Segment-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention After Clopidogrel Loading. A Randomized Double-Blind Trial J. Mehilli, A. Kastrati, S. Schulz, S. Frungel, S. G. Nekolla, W. Moshage, F. Dotzer, K. Huber, J. Pache, J. Dirschinger, M. Seyfarth, S. Martinoff, M. Schwaiger and A. Schomig *Circulation* (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19332467](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19332467)

Percutaneous coronary intervention or coronary artery bypass graft for unprotected left main coronary artery disease: the endless debate E. Meliga, M. Valgimigli, P. Buszman and P. W. Serruys *J Am Coll Cardiol* (2008) 52; 582-4; author reply 84-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18687257](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18687257)

Drug-eluting stents: a critique N. Melikian and W. Wijns *Heart* (2008) 94; 145-52

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18195121](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18195121)

Lack of benefit from percutaneous intervention of persistently occluded infarct arteries after the acute phase of myocardial infarction is time independent: insights from Occluded Artery Trial V. Menon, C. A. Pearte, C. E. Buller, P. G. Steg, S. A. Forman, H.

D. White, P. N. Marino, D. G. Katritsis, P. Caramori, R. Lasevitch, K. Loboz-Grudzien, A. Zurakowski, G. A. Lamas and J. S. Hochman *Eur Heart J* (2009) 30; 183-91

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19028780](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19028780)

The prognostic value of combined intracoronary pressure and blood flow velocity measurements after deferral of percutaneous coronary intervention M. Meuwissen, S. A. Chamuleau, M. Siebes, R. J. de Winter, K. T. Koch, L. M. Dijkstra, A. J. van den Berg, J. G. Tijssen, J. A. Spaan and J. J. Piek *Catheter Cardiovasc Interv* (2008) 71; 291-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18288725](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18288725)

Percutaneous coronary intervention utilizing a new endothelial progenitor cells antibody-coated stent: a prospective single-center registry in high-risk patients M. Miglionico, G. Patti, A. D'Ambrosio and G. Di Sciascio *Catheter Cardiovasc Interv* (2008) 71; 600-4

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18360849](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18360849)

Trifurcation left main coronary artery percutaneous revascularization--pushing catheters and envelopes D. J. Moliterno *Catheter Cardiovasc Interv* (2009) 73; 32-3

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19089933](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19089933)

Mechanical reperfusion: treat well, treat on time too G. Montalescot *Lancet* (2008) 372; 509-10

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18707968](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18707968)

Prasugrel compared with clopidogrel in patients undergoing percutaneous coronary intervention for ST-elevation myocardial infarction (TRITON-TIMI 38): double-blind, randomised controlled trial G. Montalescot, S. D. Wiviott, E. Braunwald, S. A. Murphy, C. M. Gibson, C. H. McCabe and E. M. Antman *Lancet* (2009) 373; 723-31

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19249633](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19249633)

Percutaneous revascularization of the "unprotected" left main during acute myocardial infarction: "Dare to struggle! Dare to win!" D. A. Morrison *Catheter Cardiovasc Interv* (2009) 73; 22-4

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19089931](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19089931)

The impact of gender on outcomes of patients with ST elevation myocardial infarction transported for percutaneous coronary intervention: analysis of the PRAGUE-1 and 2 studies Z. Motovska, P. Widimsky and M. Aschermann *Heart* (2008) 94; e5

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17693459](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17693459)

Effectiveness of PCI for non-acute coronary artery disease D. Mukherjee and D. J. Moliterno *Lancet* (2009) 373; 870-2

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19286067](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19286067)

Predictors and time-related impact of distal embolization during primary angioplasty M. Napodano, A. Ramondo, G. Tarantini, D. Peluso, S. Compagno, C. Fraccaro, A. C.

Frigo, R. Razzolini and S. Iliceto *Eur Heart J* (2009) 30; 305-13  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19153179](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19153179)

Periprocedural bleeding and 1-year outcome after percutaneous coronary interventions: appropriateness of including bleeding as a component of a quadruple end point G. Ndrepepa, P. B. Berger, J. Mehilli, M. Seyfarth, F. J. Neumann, A. Schomig and A. Kastrati *J Am Coll Cardiol* (2008) 51; 690-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18279731](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18279731)

One-year clinical outcomes with abciximab vs. placebo in patients with non-ST-segment elevation acute coronary syndromes undergoing percutaneous coronary intervention after pre-treatment with clopidogrel: results of the ISAR-REACT 2 randomized trial G. Ndrepepa, A. Kastrati, J. Mehilli, F. J. Neumann, J. ten Berg, O. Bruskina, F. Dotzer, M. Seyfarth, J. Pache, J. Dirschinger, P. B. Berger and A. Schomig *Eur Heart J* (2008) 29; 455-61  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18158289](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18158289)

Prognostic significance of epicardial blood flow before and after percutaneous coronary intervention in patients with acute coronary syndromes G. Ndrepepa, J. Mehilli, S. Schulz, R. Iijima, D. Keta, R. A. Byrne, J. Pache, M. Seyfarth, A. Schomig and A. Kastrati *J Am Coll Cardiol* (2008) 52; 512-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18687242](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18687242)

A prospective, double-blind, randomized, controlled trial on the efficacy and cardiorenal safety of iodixanol vs. iopromide in patients with chronic kidney disease undergoing coronary angiography with or without percutaneous coronary intervention B. Nie, W. J. Cheng, Y. F. Li, Z. Cao, Q. Yang, Y. X. Zhao, Y. H. Guo and Y. J. Zhou *Catheter Cardiovasc Interv* (2008) 72; 958-65  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19021282](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19021282)

Prognostic value of troponin after elective percutaneous coronary intervention: A meta-analysis M. B. Nienhuis, J. P. Ottervanger, H. J. Bilo, B. D. Dikkeschei and F. Zijlstra *Catheter Cardiovasc Interv* (2008) 71; 318-24  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18288753](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18288753)

Impact of platelet glycoprotein IIb/IIIa receptor inhibitors on outcomes of diabetic patients undergoing percutaneous coronary interventions using sirolimus-eluting stents E. Nikolsky, D. R. Holmes, R. Mehran, G. Dangas, E. Schampaert, M. C. Morice, J. Schofer, J. E. Sousa, M. Fahy, Y. Na, D. J. Donohoe, J. W. Moses and M. B. Leon *Catheter Cardiovasc Interv* (2008) 71; 896-906  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18498145](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18498145)

Thrombus aspiration during percutaneous coronary intervention J. Oh, N. Hong and S. M. Kang *Lancet* (2008) 372; 1034; author reply 34-5  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18805324](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18805324)

Can direct stenting in selected saphenous vein graft lesions be considered an alternative to percutaneous intervention with a distal protection device? T. Okabe, J. Lindsay, R. Torguson, D. H. Steinberg, P. Roy, T. L. Slottow, K. Kaneshige, Z. Xue, L. F. Satler, K. M. Kent, A. D. Pichard and R. Waksman *Catheter Cardiovasc Interv* (2008) 72; 799-803

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19006243](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19006243)

Long-term effectiveness of early administration of glycoprotein IIb/IIIa agents to real-world patients undergoing primary percutaneous interventions: results of a registry study in an ST-elevation myocardial infarction network P. Ortolani, A. Marzocchi, C. Marrozzini, T. Palmerini, F. Saia, N. Taglieri, F. Baldazzi, G. Dall'Ara, P. Nardini, S. Gianstefani, P. Guastaroba, R. Grilli and A. Branzi *Eur Heart J* (2009) 30; 33-43

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18996956](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18996956)

Predictive value of high sensitivity C-reactive protein in patients with ST-elevation myocardial infarction treated with percutaneous coronary intervention P. Ortolani, A. Marzocchi, C. Marrozzini, T. Palmerini, F. Saia, N. Taglieri, F. Baldazzi, S. Silenzi, M. L. Bacchi-Reggiani, P. Guastaroba, R. Grilli and A. Branzi *Eur Heart J* (2008) 29; 1241-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17766280](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17766280)

Temporal pattern of ischemic events in relation to dual antiplatelet therapy in patients with unprotected left main coronary artery stenosis undergoing percutaneous coronary intervention T. Palmerini, A. Marzocchi, C. Tamburino, I. Sheiban, M. Margheri, G. Vecchi, G. Sangiorgi, A. Santarelli, A. L. Bartorelli, C. Briguori, L. Vignali, F. Di Pede, A. Ramondo, L. Inglese, M. De Carlo, L. Bolognese, A. Benassi, C. Palmieri, V. Filippone, D. Sangiorgi, F. Barlocco, G. Lauria and S. De Servi *J Am Coll Cardiol* (2009) 53; 1176-81

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19341857](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19341857)

Long-term mortality after percutaneous coronary intervention with drug-eluting stent implantation versus coronary artery bypass surgery for the treatment of multivessel coronary artery disease D. W. Park, S. C. Yun, S. W. Lee, Y. H. Kim, C. W. Lee, M. K. Hong, J. J. Kim, S. J. Choo, H. Song, C. H. Chung, J. W. Lee, S. W. Park and S. J. Park *Circulation* (2008) 117; 2079-86

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18413495](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18413495)

Current status of percutaneous coronary intervention with drug-eluting stents in Asia S. J. Park and Y. H. Kim *Circulation* (2008) 118; 2730-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19106397](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19106397)

Point-of-care measurement of clopidogrel responsiveness predicts clinical outcome in patients undergoing percutaneous coronary intervention results of the ARMYDA-PRO (Antiplatelet therapy for Reduction of MYocardial Damage during Angioplasty-Platelet Reactivity Predicts Outcome) study G. Patti, A. Nusca, F. Mangiacapra, L. Gatto, A. D'Ambrosio and G. Di Sciascio *J Am Coll Cardiol* (2008) 52; 1128-33

<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list\_uids=18804738

Outcome in transferred and nontransferred patients after primary percutaneous coronary intervention for ischaemic out-of-hospital cardiac arrest H. O. Peels, G. A. Jessurun, I. C. van der Horst, A. E. Arnold, L. H. Piers and F. Zijlstra *Catheter Cardiovasc Interv* (2008) 71; 147-51

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18231992](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18231992)

Effects of off-pump versus on-pump coronary artery bypass grafting on early and late right ventricular function T. J. Pegg, J. B. Selvanayagam, T. D. Karamitsos, R. J. Arnold, J. M. Francis, S. Neubauer and D. P. Taggart *Circulation* (2008) 117; 2202-10

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18413501](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18413501)

Comparison of five-year outcome in African Americans versus Caucasians following percutaneous coronary intervention J. Pradhan, T. L. Schreiber, A. Niraj, V. Veeranna, K. Ramesh, L. Saigh and L. Afonso *Catheter Cardiovasc Interv* (2008) 72; 36-44

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18383170](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18383170)

Primary percutaneous coronary intervention for acute myocardial infarction caused by unprotected left main stem thrombosis S. B. Prasad, R. Whitbourn, Y. Malaiapan, W. Ahmar, A. MacIsaac and I. T. Meredith *Catheter Cardiovasc Interv* (2009) 73; 301-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19213083](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19213083)

Cardiac optical coherence tomography O. C. Raffel, T. Akasaka and I. K. Jang *Heart* (2008) 94; 1200-10

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18703696](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18703696)

In vivo association between positive coronary artery remodelling and coronary plaque characteristics assessed by intravascular optical coherence tomography O. C. Raffel, F. M. Merchant, G. J. Tearney, S. Chia, D. D. Gauthier, E. Pomerantsev, K. Mizuno, B. E. Bouma and I. K. Jang *Eur Heart J* (2008) 29; 1721-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18577556](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18577556)

Integration of multislice computed tomography with magnetic navigation facilitates percutaneous coronary interventions without additional contrast agents S. Ramcharitar, F. Pugliese, C. Schultz, J. Ligthart, P. de Feyter, H. Li, N. Mollet, M. van de Ent, P. W. Serruys and R. J. van Geuns *J Am Coll Cardiol* (2009) 53; 741-6

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19245963](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19245963)

A randomized comparison of the magnetic navigation system versus conventional percutaneous coronary intervention S. Ramcharitar, R. J. van Geuns, M. Patterson, W. J. van der Giessen, M. van der Ent, R. T. van Domburg and P. W. Serruys *Catheter Cardiovasc Interv* (2008) 72; 761-70

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18798236](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18798236)

Electrophysiological effects of late percutaneous coronary intervention for infarct-related coronary artery occlusion: the Occluded Artery Trial-Electrophysiological Mechanisms



(OAT-EP) E. J. Rashba, G. A. Lamas, J. P. Couderc, S. M. Hollist, V. Dzavik, W. Ruzylo, V. Fridrich, C. E. Buller, S. A. Forman, J. A. Kufera, A. C. Carvalho and J. S. Hochman *Circulation* (2009) 119; 779-87  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19188505](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19188505)

Coronary thrombectomy by retrieval of an open emboli-protection filter device I. Reho, C. Gruner and M. Roffi *Heart* (2008) 94; 274  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18276814](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18276814)

Immediate versus deferred coronary angioplasty in non-ST-elevation acute coronary syndromes R. K. Riezebos, E. Ronner, E. Ter Bals, T. Slagboom, P. C. Smits, J. M. Ten Berg, F. Kiemeneij, G. Amoroso, M. S. Patterson, M. J. Suttorp, J. G. Tijssen and G. J. Laarman *Heart* (2008)  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19098058](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19098058)

Nonrandomized comparison of coronary artery bypass surgery and percutaneous coronary intervention for the treatment of unprotected left main coronary artery disease in octogenarians J. Rodes-Cabau, J. Deblois, O. F. Bertrand, S. Mohammadi, J. Courtis, E. Larose, F. Dagenais, J. P. Dery, P. Mathieu, M. Rousseau, G. Barbeau, R. Baillot, O. Gleeton, J. Perron, C. M. Nguyen, L. Roy, D. Doyle, R. De Larochelliere, P. Bogaty and P. Voisine *Circulation* (2008) 118; 2374-81  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19029471](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19029471)

Long-term safety and durability of percutaneous septal sinus shortening (The PS(3) System) in an ovine model J. H. Rogers, D. A. Rahdert, G. R. Caputo, P. A. Takeda, I. F. Palacios, F. O. Tio, E. A. Taylor and R. I. Low *Catheter Cardiovasc Interv* (2009) 73; 540-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19235241](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19235241)

Thrombus aspiration during primary percutaneous coronary intervention improves myocardial reperfusion and reduces infarct size: the EXPIRA (thrombectomy with export catheter in infarct-related artery during primary percutaneous coronary intervention) prospective, randomized trial G. Sardella, M. Mancone, C. Bucciarelli-Ducci, L. Agati, R. Scardala, I. Carbone, M. Francone, A. Di Roma, G. Benedetti, G. Conti and F. Fedele *J Am Coll Cardiol* (2009) 53; 309-15  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19161878](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19161878)

A meta-analysis of 17 randomized trials of a percutaneous coronary intervention-based strategy in patients with stable coronary artery disease A. Schomig, J. Mehilli, A. de Waha, M. Seyfarth, J. Pache and A. Kastrati *J Am Coll Cardiol* (2008) 52; 894-904  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18772058](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18772058)

Optimal medical therapy with or without percutaneous coronary intervention to reduce ischemic burden: results from the Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation (COURAGE) trial nuclear substudy L. J. Shaw, D. S. Berman, D. J. Maron, G. B. Mancini, S. W. Hayes, P. M. Hartigan, W. S. Weintraub, R.

A. O'Rourke, M. Dada, J. A. Spertus, B. R. Chaitman, J. Friedman, P. Slomka, G. V. Heller, G. Germano, G. Gosselin, P. Berger, W. J. Kostuk, R. G. Schwartz, M. Knudtson, E. Veledar, E. R. Bates, B. McCallister, K. K. Teo and W. E. Boden *Circulation* (2008) 117; 1283-91

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18268144](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18268144)

Early and long-term results of percutaneous coronary intervention for unprotected left main trifurcation disease I. Sheiban, A. Gerasimou, M. Bollati, G. Biondi-Zoccai, F. Sciuto, P. Omede, D. Sillano, G. P. Trevi and C. Moretti *Catheter Cardiovasc Interv* (2009) 73; 25-31

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19089932](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19089932)

Impact of bivalirudin or unfractionated heparin on platelet aggregation in patients pretreated with 600 mg clopidogrel undergoing elective percutaneous coronary intervention D. Sibbing, G. Busch, S. Braun, S. Jawansky, A. Schomig, A. Kastrati, I. Ott and N. von Beckerath *Eur Heart J* (2008) 29; 1504-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18467322](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18467322)

Cytochrome P450 2C19 loss-of-function polymorphism and stent thrombosis following percutaneous coronary intervention D. Sibbing, J. Stegherr, W. Latz, W. Koch, J. Mehilli, K. Dorrlor, T. Morath, A. Schomig, A. Kastrati and N. von Beckerath *Eur Heart J* (2009)

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19193675](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19193675)

Genetic determinants of response to clopidogrel and cardiovascular events T. Simon, C. Verstuyft, M. Mary-Krause, L. Quteineh, E. Drouet, N. Meneveau, P. G. Steg, J. Ferrieres, N. Danchin and L. Becquemont *N Engl J Med* (2009) 360; 363-75

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19106083](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19106083)

Mayo Clinic Risk Score for percutaneous coronary intervention predicts in-hospital mortality in patients undergoing coronary artery bypass graft surgery M. Singh, B. J. Gersh, S. Li, J. S. Rumsfeld, J. A. Spertus, S. M. O'Brien, R. M. Suri and E. D. Peterson *Circulation* (2008) 117; 356-62

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18172033](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18172033)

Mortality differences between men and women after percutaneous coronary interventions. A 25-year, single-center experience M. Singh, C. S. Rihal, B. J. Gersh, V. L. Roger, M. R. Bell, R. J. Lennon, A. Lerman and D. R. Holmes, Jr. *J Am Coll Cardiol* (2008) 51; 2313-20

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18549915](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18549915)

Comorbid conditions and outcomes after percutaneous coronary intervention M. Singh, C. S. Rihal, V. L. Roger, R. J. Lennon, J. Spertus, A. Jahangir and D. R. Holmes, Jr. *Heart* (2008) 94; 1424-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17923464](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17923464)

Surgery, not percutaneous revascularization, is the preferred strategy for patients with

significant left main coronary stenosis C. R. Smith *Circulation* (2009) 119; 1013-20  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19237672](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19237672)

Coronary stents and chronic anticoagulation A. Sourgounis, J. Lipiecki, T. S. Lo and M. Hamon *Circulation* (2009) 119; 1682-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19332481](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19332481)

Acute vessel occlusion after coronary stenting: intravascular ultrasound diagnosis of extensive intramural haematoma G. Souteyrand and P. Motreff *Eur Heart J* (2008) 29; 9  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17675314](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17675314)

Adjunctive thrombectomy and distal protection in primary percutaneous coronary intervention: impact on microvascular perfusion and outcomes M. Srinivasan, C. Rihal, D. R. Holmes and A. Prasad *Circulation* (2009) 119; 1311-9  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19273732](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19273732)

The CIAO (Coronary Interventions Antiplatelet-based Only) Study: a randomized study comparing standard anticoagulation regimen to absence of anticoagulation for elective percutaneous coronary intervention E. Stabile, W. Nammias, L. Salemme, G. Sorropago, A. Cioppa, T. Tesorio, V. Ambrosini, E. Campopiano, G. Popusoi, G. Biondi Zoccai and P. Rubino *J Am Coll Cardiol* (2008) 52; 1293-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18929239](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18929239)

Aspirin following PCI: too much of a good thing? S. R. Steinhubl and P. B. Berger *Eur Heart J* (2009)  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19293194](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19293194)

Aspiration of embolized thrombus during primary percutaneous coronary intervention M. G. Stoel, C. von Birgelen and F. Zijlstra *Catheter Cardiovasc Interv* (2008)  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19180655](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19180655)

Angioplasty strategies in ST-segment-elevation myocardial infarction: part I: primary percutaneous coronary intervention G. W. Stone *Circulation* (2008) 118; 538-51  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18663102](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18663102)

Methods to reduce patients' maximum skin dose during percutaneous coronary intervention for chronic total occlusion S. Suzuki, S. Furui, T. Isshiki, K. Kozuma, Y. Koyama, M. Ochiai and Y. Asakura *Catheter Cardiovasc Interv* (2008) 71; 792-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18324699](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18324699)

Patients' skin dose during percutaneous coronary intervention for chronic total occlusion S. Suzuki, S. Furui, T. Isshiki, K. Kozuma, Y. Koyama, H. Yamamoto, M. Ochiai, Y. Asakura and Y. Ikari *Catheter Cardiovasc Interv* (2008) 71; 160-4  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17932885](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17932885)

Primary percutaneous balloon pericardiotomy for malignant pericardial effusion N.

Swanson, I. Mirza, N. Wijesinghe and G. Devlin Catheter Cardiovasc Interv (2008) 71; 504-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18307242](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18307242)

Does prior PCI increase the risk of subsequent CABG? D. P. Taggart Eur Heart J (2008) 29; 573-5  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18285359](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18285359)

Percutaneous coronary intervention using a novel 4-French coronary accessor S. Takeshita, T. Shiono, A. Takagi, T. Ito and S. Saito Catheter Cardiovasc Interv (2008) 72; 222-7  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18546232](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18546232)

Complete versus incomplete revascularization in patients with multivessel disease undergoing percutaneous coronary intervention with drug-eluting stents C. Tamburino, D. J. Angiolillo, P. Capranzano, K. Dimopoulos, A. La Manna, R. Barbagallo, F. Tagliareni, S. Mangiafico, L. A. Guzman, A. R. Galassi and T. A. Bass Catheter Cardiovasc Interv (2008) 72; 448-56  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18814218](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18814218)

Percutaneous revascularization is the preferred strategy for patients with significant left main coronary stenosis P. S. Teirstein Circulation (2009) 119; 1021-33  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19237673](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19237673)

Value of ST-resolution analysis in the era of primary percutaneous coronary intervention C. J. Terkelsen and H. R. Andersen Heart (2008) 94; 13-5  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18083948](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18083948)

Percutaneous coronary intervention related delay in ST-segment elevation myocardial infarction patients C. J. Terkelsen, J. T. Sorensen, T. T. Nielsen and E. Boersma J Am Coll Cardiol (2009) 53; 1244; author reply 44-5  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19341870](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19341870)

Intracoronary compared with intravenous bolus abciximab application in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention: the randomized Leipzig immediate percutaneous coronary intervention abciximab IV versus IC in ST-elevation myocardial infarction trial H. Thiele, K. Schindler, J. Friedenberger, I. Eitel, G. Furnau, E. Grebe, S. Erbs, A. Linke, S. Mobius-Winkler, D. Kivelitz and G. Schuler Circulation (2008) 118; 49-57  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18559698](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18559698)

Local vessel injury following percutaneous coronary intervention does not promote early mobilisation of endothelial progenitor cells in the absence of myocardial necrosis H. E. Thomas, P. J. Avery, J. M. Ahmed, R. Edwards, I. Purcell, A. G. Zaman, H. M. Arthur and B. D. Keavney Heart (2009) 95; 555-8  
<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list\_uids=18977801

Cytochrome P450 2C19 681G>A polymorphism and high on-clopidogrel platelet reactivity associated with adverse 1-year clinical outcome of elective percutaneous coronary intervention with drug-eluting or bare-metal stents D. Trenk, W. Hochholzer, M. F. Fromm, L. E. Chialda, A. Pahl, C. M. Valina, C. Stratz, P. Schmiebusch, H. P. Bestehorn, H. J. Buttner and F. J. Neumann *J Am Coll Cardiol* (2008) 51; 1925-34  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18482659](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18482659)

Percutaneous coronary interventions for non-acute coronary artery disease: a quantitative 20-year synopsis and a network meta-analysis T. A. Trikalinos, A. A. Alsheikh-Ali, A. Tatsioni, B. K. Nallamothu and D. M. Kent *Lancet* (2009) 373; 911-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19286090](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19286090)

The correlation between lipid volume in the target lesion, measured by integrated backscatter intravascular ultrasound, and post-procedural myocardial infarction in patients with elective stent implantation T. Uetani, T. Amano, H. Ando, K. Yokoi, K. Arai, M. Kato, N. Marui, M. Nanki, T. Matsubara, H. Ishii, H. Izawa and T. Murohara *Eur Heart J* (2008) 29; 1714-20  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18559327](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18559327)

Distal myocardial protection with intracoronary beta blocker when added to a Gp IIb/IIIa platelet receptor blocker during percutaneous coronary intervention improves clinical outcome B. F. Uretsky, Y. Birnbaum, A. Osman, R. Gupta, O. Paniagua, A. Chamoun, A. Pohwani, C. Lui, E. Lev, T. McGehee, D. Kumar, A. Akhtar, A. Anzuini, E. R. Schwarz and F. W. Wang *Catheter Cardiovasc Interv* (2008) 72; 488-97  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18814223](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18814223)

Percutaneous coronary intervention for a right coronary artery stent occlusion using retrograde delivery of a sirolimus-eluting stent via a septal perforator M. Utunomiya, O. Katoh and S. Nakamura *Catheter Cardiovasc Interv* (2009) 73; 475-80  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19229979](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19229979)

Percutaneous coronary interventions in octogenarians: Acute and 12 month results in a large single-centre experience E. Varani, M. Aquilina, M. Balducelli, G. Vecchi, V. Frassinetti and A. Maresta *Catheter Cardiovasc Interv* (2009) 73; 449-54  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19229976](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19229976)

Single or multivessel percutaneous coronary intervention in ST-elevation myocardial infarction patients E. Varani, M. Balducelli, M. Aquilina, G. Vecchi, M. N. Hussien, V. Frassinetti and A. Maresta *Catheter Cardiovasc Interv* (2008) 72; 927-33  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18798239](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18798239)

Impact of pretreatment with clopidogrel on initial patency and outcome in patients treated with primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: a systematic review P. J. Vlaar, T. Svilaas, K. Damman, B. J. de Smet, J. G. Tijssen, H. L. Hillege and F. Zijlstra *Circulation* (2008) 118; 1828-36

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18852370](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18852370)

Cardiac death and reinfarction after 1 year in the Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction Study (TAPAS): a 1-year follow-up study P. J. Vlaar, T. Svilaas, I. C. van der Horst, G. F. Diercks, M. L. Fokkema, B. J. de Smet, A. F. van den Heuvel, R. L. Anthonio, G. A. Jessurun, E. S. Tan, A. J. Suurmeijer and F. Zijlstra *Lancet* (2008) 371; 1915-20

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18539223](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18539223)

Computer-assisted myocardial blush quantification after percutaneous coronary angioplasty for acute myocardial infarction: a substudy from the TAPAS trial M. Vogelzang, P. J. Vlaar, T. Svilaas, D. Amo, M. W. Nijsten and F. Zijlstra *Eur Heart J* (2009) 30; 594-9

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19168868](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19168868)

Unusual use of a standard percutaneous transluminal coronary angioplasty balloon catheter to the rescue M. Vorpahl, K. L. Laugwitz, B. Zrenner, A. Kastrati and A. Schomig *Eur Heart J* (2008) 29; 2732

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18515804](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18515804)

Circulating lipid hydroperoxides predict cardiovascular events in patients with stable coronary artery disease: the PREVENT study M. F. Walter, R. F. Jacob, R. E. Bjork, B. Jeffers, J. Buch, Y. Mizuno and R. P. Mason *J Am Coll Cardiol* (2008) 51; 1196-202

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18355658](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18355658)

Patterns of cardiac marker surveillance after elective percutaneous coronary intervention and implications for the use of periprocedural myocardial infarction as a quality metric: a report from the National Cardiovascular Data Registry (NCDR) T. Y. Wang, E. D. Peterson, D. Dai, H. V. Anderson, S. V. Rao, R. G. Brindis and M. T. Roe *J Am Coll Cardiol* (2008) 51; 2068-74

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18498965](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18498965)

Acute myocardial infarction H. D. White and D. P. Chew *Lancet* (2008) 372; 570-84

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18707987](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18707987)

Safety and efficacy of bivalirudin with and without glycoprotein IIb/IIIa inhibitors in patients with acute coronary syndromes undergoing percutaneous coronary intervention 1-year results from the ACUITY (Acute Catheterization and Urgent Intervention Triage strategY) trial H. D. White, E. M. Ohman, A. M. Lincoff, M. E. Bertrand, A. Colombo, B. T. McLaurin, D. A. Cox, S. J. Pocock, J. A. Ware, S. V. Manoukian, A. J. Lansky, R. Mehran, J. W. Moses and G. W. Stone *J Am Coll Cardiol* (2008) 52; 807-14

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18755342](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18755342)

Routine use of oxygen in the treatment of myocardial infarction: systematic review M. Wijesinghe, K. Perrin, A. Ranchord, M. Simmonds, M. Weatherall and R. Beasley *Heart* (2009) 95; 198-202

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18708420](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18708420)

Intensive oral antiplatelet therapy for reduction of ischaemic events including stent thrombosis in patients with acute coronary syndromes treated with percutaneous coronary intervention and stenting in the TRITON-TIMI 38 trial: a subanalysis of a randomised trial S. D. Wiviott, E. Braunwald, C. H. McCabe, I. Horvath, M. Keltai, J. P. Herrman, F. Van de Werf, W. E. Downey, B. M. Scirica, S. A. Murphy and E. M. Antman *Lancet* (2008) 371; 1353-63

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18377975](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18377975)

Determinants of thrombin generation, fibrinolytic activity, and endothelial dysfunction in patients on dual antiplatelet therapy: involvement of factors other than platelet aggregability in Virchow's triad Y. Yano, T. Ohmori, S. Hoshida, S. Madoiwa, K. Yamamoto, T. Katsuki, T. Mitsuhashi, J. Mimuro, K. Shimada, K. Kario and Y. Sakata *Eur Heart J* (2008) 29; 1729-38

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18270211](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18270211)

Volume-outcome relation for contemporary percutaneous coronary interventions (PCI) in daily clinical practice: is it limited to high-risk patients? Results from the Registry of Percutaneous Coronary Interventions of the Arbeitsgemeinschaft Leitende Kardiologische Krankenhausärzte (ALKK) R. Zahn, M. Gottwik, M. Hochadel, J. Senges, U. Zeymer, A. Vogt, T. Meinertz, R. Dietz, K. E. Hauptmann, E. Grube, S. Kerber and U. Sechtem *Heart* (2008) 94; 329-35

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17664190](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17664190)