

Duration of symptoms is the key modulator of the choice of reperfusion for ST-elevation myocardial infarction P. W. Armstrong, C. M. Westerhout and R. C. Welsh Circulation (2009) 119; 1293-303

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19273730](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19273730)

Late target lesion revascularization after implantation of sirolimus-eluting stent M. K. Hong, G. S. Mintz, C. W. Lee, D. W. Park, S. W. Lee, Y. H. Kim, I. H. Jung, S. H. Kim, S. S. Cheong, J. J. Kim, S. W. Park and S. J. Park Catheter Cardiovasc Interv (2008) 71; 299-303

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=17985380](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=17985380)

Beyond established and novel risk factors: lifestyle risk factors for cardiovascular disease D. Mozaffarian, P. W. Wilson and W. B. Kannel Circulation (2008) 117; 3031-8  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18541753](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18541753)

Nuclear factor-{kappa}B activation contributes to vascular endothelial dysfunction via oxidative stress in overweight/obese middle-aged and older humans G. L. Pierce, L. A. Lesniewski, B. R. Lawson, S. D. Beske and D. R. Seals Circulation (2009) 119; 1284-92  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19237660](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19237660)

Association of blood glucose with angiographic and clinical outcomes among patients with ST-segment elevation myocardial infarction (from the CLARITY-TIMI-28 study) D. S. Pinto, A. J. Kirtane, Y. B. Pride, S. A. Murphy, M. S. Sabatine, C. P. Cannon and C. M. Gibson Am J Cardiol (2008) 101; 303-7

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18237589](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18237589)

Observations and outcomes of definite and probable drug-eluting stent thrombosis seen at a single hospital in a four-year period T. L. Pinto Slottow, D. H. Steinberg, P. K. Roy, A. N. Buch, T. Okabe, Z. Xue, K. Kaneshige, R. Torguson, J. Lindsay, A. D. Pichard, L. F. Satler, W. O. Suddath, K. M. Kent and R. Waksman Am J Cardiol (2008) 102; 298-303

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18638590](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18638590)

Carotid artery stent placement is safe in the very elderly (> or =80 years) C. A. Velez, C. J. White, J. P. Reilly, J. S. Jenkins, T. J. Collins, M. A. Grise, P. W. McMullan and S. R. Ramee Catheter Cardiovasc Interv (2008) 72; 303-8

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18726941](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18726941)