

Usefulness of coronary fractional flow reserve measurements in guiding clinical decisions in intermediate or equivocal left main coronary stenoses J. Courtis, J. Rodes-Cabau, E. Larose, J. M. Potvin, J. P. Dery, R. D. Larocheilliere, M. Cote, O. Counterousse, C. M. Nguyen, G. Proulx, S. Rinfret and O. F. Bertrand Am J Cardiol (2009) 103; 943-9  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19327420](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19327420)

Endovascular repair of abdominal aortic aneurysm R. M. Greenhalgh and J. T. Powell N Engl J Med (2008) 358; 494-501  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18234753](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18234753)

Comparison of coronary flow reserve and fractional flow reserve in patients with versus without diabetes mellitus and having elective percutaneous coronary intervention and abciximab therapy (from the PREDICT Trial) A. S. Kini, M. C. Kim, P. R. Moreno, P. Krishnan, O. C. Ivan and S. K. Sharma Am J Cardiol (2008) 101; 796-800  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18328843](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18328843)

An integrated TAXUS IV, V, and VI intravascular ultrasound analysis of the predictors of edge restenosis after bare metal or paclitaxel-eluting stents J. Liu, A. Maehara, G. S. Mintz, N. J. Weissman, A. Yu, H. Wang, L. Mandinov, J. J. Popma, S. G. Ellis, E. Grube, K. D. Dawkins and G. W. Stone Am J Cardiol (2009) 103; 501-6  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19195510](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195510)