

Long-term clinical benefit of sirolimus-eluting stents in patients with in-stent restenosis results of the RIBS-II (Restenosis Intra-stent: Balloon angioplasty vs. elective sirolimus-eluting Stenting) study F. Alfonso, M. J. Perez-Vizcayno, R. Hernandez, A. Bethencourt, V. Marti, J. R. Lopez-Minguez, J. Angel, A. Iniguez, C. Moris, A. Cequier, M. Sabate, J. Escaned, P. Jimenez-Quevedo, C. Banuelos, A. Suarez and C. Macaya J Am Coll Cardiol (2008) 52; 1621-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18992651

Sirolimus-eluting stents versus bare-metal stents in patients with in-stent restenosis: results of a pooled analysis of two randomized studies F. Alfonso, M. J. Perez-Vizcayno, R. Hernandez, C. Fernandez, J. Escaned, C. Banuelos, A. Bethencourt, J. R. Lopez-Minguez, J. Angel, A. Cequier, M. Sabate, C. Moris, J. Zueco and R. Seabra-Gomes Catheter Cardiovasc Interv (2008) 72; 459-67

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18814274

Treatment of drug-eluting stent restenosis: An emerging challenge A. Aminian, T. Kabir and E. Eeckhout Catheter Cardiovasc Interv (2008)

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19199365

Tailored clopidogrel loading dose according to platelet reactivity monitoring to prevent acute and subacute stent thrombosis L. Bonello, L. Camoin-Jau, S. Armero, O. Com, S. Arques, C. Burignat-Bonello, M. P. Giacomoni, R. Bonello, F. Collet, P. Rossi, P. Barragan, F. Dignat-George and F. Paganelli Am J Cardiol (2009) 103; 5-10

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19101221

Comparison of outcomes of drug-eluting stents versus bare-metal stents in nonostial proximal left anterior descending coronary arteries L. Bonello, A. De Labriolle, G. Lemesle, P. Roy, D. H. Steinberg, T. L. Slottow, Z. Xue, R. Torguson, K. Kaneshige, W. O. Suddath, L. F. Satler, K. M. Kent, J. Lindsay, A. D. Pichard and R. Waksman Am J Cardiol (2009) 103; 496-500

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195509

Four-year clinical follow-up of the rapamycin-eluting stent evaluated at Rotterdam Cardiology Hospital registry J. Daemen, N. Kukreja, P. H. van Twisk, Y. Onuma, P. P. de Jaegere, R. van Domburg and P. W. Serruys Am J Cardiol (2008) 101; 1105-11
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18394442

Predictive factors for in-stent late loss and coronary lesion progression in patients with type 2 diabetes mellitus randomized to rosiglitazone or placebo A. V. Finn, J. S. Oh, M. Hendricks, M. Daher, E. Cagliero, R. M. Byrne, J. Nadelson, J. Crimins, A. Kastrati, A. Schomig, O. Bruskina, I. Palacios, M. C. John and H. K. Gold Am Heart J (2009) 157; 383 e1-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19185649

Contribution of organized thrombus to in-stent restenosis after sirolimus-eluting stent implantation: optical coherence tomography findings K. Fujii, M. Masutani and M.

Ohyanagi Eur Heart J (2008) 29; 1385

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18077442

Balancing the risks of restenosis and stent thrombosis in bare-metal versus drug-eluting stents: results of a decision analytic model P. Garg, D. J. Cohen, T. Gaziano and L. Mauri J Am Coll Cardiol (2008) 51; 1844-53

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18466798

Comparative safety and efficacy of a sirolimus-eluting versus paclitaxel-eluting stent: a meta-analysis H. S. Gurm, T. Boyden and K. B. Welch Am Heart J (2008) 155; 630-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18371469

Late drug-eluting stent thrombosis of unprotected left main coronary artery lesions--sometimes possible but rarely definite or probable H. Hamdalla and D. J. Moliterno Eur Heart J (2008)

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18664463

Late stent malapposition risk is higher after drug-eluting stent compared with bare-metal stent implantation and associates with late stent thrombosis A. K. Hassan, S. C.

Bergheanu, T. Stijnen, B. L. van der Hoeven, J. D. Snoep, J. W. Plevier, M. J. Schalij and J. W. Jukema Eur Heart J (2009)

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19158118

Usefulness of 64-detector computed tomographic angiography for diagnosing in-stent restenosis in native coronary arteries H. S. Hecht, M. Zaric, V. Jelnin, L. Lubarsky, M. Prakash and G. Roubin Am J Cardiol (2008) 101; 820-4

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18328847

Clinical presentation of patients with late target lesion revascularization H. Kaneda Catheter Cardiovasc Interv (2008) 72; 749

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18942148

A novel bioresorbable polymer paclitaxel-eluting stent for the treatment of single and multivessel coronary disease: primary results of the COSTAR (Cobalt Chromium Stent With Antiproliferative for Restenosis) II study M. W. Krucoff, D. J. Kereiakes, J. L. Petersen, R. Mehran, V. Hasselblad, A. J. Lansky, P. J. Fitzgerald, J. Garg, M. A. Turco, C. A. Simonton, 3rd, S. Verheye, C. L. Dubois, R. Gammon, W. B. Batchelor, C. D. O'Shaughnessy, J. B. Hermiller, Jr., J. Schofer, M. Buchbinder and W. Wijns J Am Coll Cardiol (2008) 51; 1543-52

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18420096

Stent strut fracture-induced restenosis in the right coronary artery: detection by MDCT S.

U. Kwon, J. H. Doh, J. Namgung and S. Y. Lee Heart (2008) 94; 221

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18195125

Drug-eluting stenting followed by cilostazol treatment reduces late restenosis in patients

with diabetes mellitus the DECLARE-DIABETES Trial (A Randomized Comparison of Triple Antiplatelet Therapy with Dual Antiplatelet Therapy After Drug-Eluting Stent Implantation in Diabetic Patients) S. W. Lee, S. W. Park, Y. H. Kim, S. C. Yun, D. W. Park, C. W. Lee, M. K. Hong, H. S. Kim, J. K. Ko, J. H. Park, J. H. Lee, S. W. Choi, I. W. Seong, Y. H. Cho, N. H. Lee, J. H. Kim, K. J. Chun and S. J. Park J Am Coll Cardiol (2008) 51; 1181-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18355656

An integrated TAXUS IV, V, and VI intravascular ultrasound analysis of the predictors of edge restenosis after bare metal or paclitaxel-eluting stents J. Liu, A. Maehara, G. S. Mintz, N. J. Weissman, A. Yu, H. Wang, L. Mandinov, J. J. Popma, S. G. Ellis, E. Grube, K. D. Dawkins and G. W. Stone Am J Cardiol (2009) 103; 501-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195510

Comparison of the sirolimus-eluting versus paclitaxel-eluting coronary stent in patients with diabetes mellitus: the diabetes and drug-eluting stent (DiabeDES) randomized angiography trial M. Maeng, L. O. Jensen, A. M. Galloe, P. Thayssen, E. H. Christiansen, K. N. Hansen, S. Helqvist, H. E. Botker, J. F. Lassen and L. Thuesen Am J Cardiol (2009) 103; 345-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19166687

Usefulness of 64-detector row computed tomography for evaluation of intracoronary stents in symptomatic patients with suspected in-stent restenosis N. Manghat, R. Van Lingen, P. Hewson, F. Syed, N. Kakani, I. Cox, C. Roobottom and G. Morgan-Hughes Am J Cardiol (2008) 101; 1567-73
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18489934

Randomized trial of three rapamycin-eluting stents with different coating strategies for the reduction of coronary restenosis J. Mehilli, R. A. Byrne, A. Wieczorek, R. Iijima, S. Schulz, O. Bruskina, J. Pache, R. Wessely, A. Schomig and A. Kastrati Eur Heart J (2008) 29; 1975-82
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18550554

Optimal treatment for in-stent restenosis after BMS--DES, coated balloon, or scalpel? D. Mukherjee Eur Heart J (2008) 29; 1595-6
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18556717

Commentary on drug-eluting stents: to use or not to use--that is the question! M. E. Pfisterer Heart (2008) 94; 1648-50
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18782796

Noninvasive assessment of coronary in-stent restenosis by dual-source computed tomography T. Pfleiderer, M. Marwan, A. Renz, S. Bachmann, D. Ropers, A. Kuettner, K. Anders, F. Bamberg, W. G. Daniel and S. Achenbach Am J Cardiol (2009) 103; 812-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19268737

Observations and outcomes of definite and probable drug-eluting stent thrombosis seen at a single hospital in a four-year period T. L. Pinto Slottow, D. H. Steinberg, P. K. Roy, A. N. Buch, T. Okabe, Z. Xue, K. Kaneshige, R. Torguson, J. Lindsay, A. D. Pichard, L. F. Satler, W. O. Suddath, K. M. Kent and R. Waksman Am J Cardiol (2008) 102; 298-303

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18638590

Commentary on drug-eluting stents: NICE technology appraisal guidance J. Richardson, A. Stevens, D. Barnett and C. Longson Heart (2008) 94; 1650-2

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18768567

Stent fracture, an incidental finding or a significant marker of clinical in-stent restenosis? F. Shaikh, R. Maddikunta, M. Djelmami-Hani, J. Solis, S. Allaqaband and T. Bajwa Catheter Cardiovasc Interv (2008) 71; 614-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18360853

Angiographic patterns of drug-eluting stent restenosis and one-year outcomes after treatment with repeated percutaneous coronary intervention E. Solinas, G. Dangas, A. J. Kirtane, A. J. Lansky, T. Franklin-Bond, P. Boland, G. Syros, Y. H. Kim, A. Gupta, G. Mintz, M. Fahy, M. Collins, S. Kodali, G. W. Stone, J. W. Moses, M. B. Leon and R. Mehran Am J Cardiol (2008) 102; 311-5

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18638592

Outcome differences with the use of drug-eluting stents for the treatment of in-stent restenosis of bare-metal stents versus drug-eluting stents D. H. Steinberg, M. A. Gaglia, Jr., T. L. Pinto Slottow, P. Roy, L. Bonello, A. De Labriolle, G. Lemesle, R. Torguson, K. Kaneshige, Z. Xue, W. O. Suddath, K. M. Kent, L. F. Satler, A. D. Pichard, J. Lindsay and R. Waksman Am J Cardiol (2009) 103; 491-5

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19195508

Long-term clinical outcomes after sirolimus-eluting stent implantation for treatment of restenosis within bare-metal versus drug-eluting stents C. Whan Lee, S. H. Kim, J. Suh, D. W. Park, S. H. Lee, Y. H. Kim, M. K. Hong, J. J. Kim, S. W. Park and S. J. Park Catheter Cardiovasc Interv (2008) 71; 594-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18311841

Rates of stent thrombosis in bare-metal versus drug-eluting stents (from a large Australian multicenter registry) B. P. Yan, S. J. Duffy, D. J. Clark, J. Lefkovits, R. Warren, R. Gurvitch, R. Lew, M. Sebastian, A. Brennan, N. Andrianopoulos, C. M. Reid and A. E. Ajani Am J Cardiol (2008) 101; 1716-22

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18549846

Macrophage migration inhibitory factor in cardiovascular disease A. Zernecke, J. Bernhagen and C. Weber Circulation (2008) 117; 1594-602

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18362243