

Comparison of assessment of native coronary arteries by standard versus three-dimensional coronary angiography P. Agostoni, G. Biondi-Zoccai, G. Van Langenhove, K. Cornelis, P. Vermeersch, C. Convens, C. Vassanelli, P. Van Den Heuvel, F. Van Den Branden and S. Verheye *Am J Cardiol* (2008) 102; 272-9
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18638585

Presence and extent of coronary artery disease by cardiac computed tomography and risk for acute coronary syndrome in cocaine users among patients with chest pain F. Bamberg, C. L. Schlett, Q. A. Truong, I. S. Rogers, W. Koenig, J. T. Nagurny, S. Seneviratne, S. J. Lehman, R. C. Cury, S. Abbara, J. Butler, H. Lee, T. J. Brady and U. Hoffmann *Am J Cardiol* (2009) 103; 620-5
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19231323

Predictive value of coronary calcifications for future cardiac events in asymptomatic individuals A. Becker, A. Leber, C. Becker and A. Knez *Am Heart J* (2008) 155; 154-60
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18082507

Usefulness of 64-slice multidetector computed tomography as an initial diagnostic approach in patients with acute chest pain S. A. Chang, S. I. Choi, E. K. Choi, H. K. Kim, J. W. Jung, E. J. Chun, K. S. Kim, Y. S. Cho, W. Y. Chung, T. J. Youn, I. H. Chae, D. J. Choi and H. J. Chang *Am Heart J* (2008) 156; 375-83
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18657674

Comparison of 64-slice multidetector computed tomography with spectral analysis of intravascular ultrasound backscatter signals for characterizations of noncalcified coronary arterial plaques B. J. Choi, D. K. Kang, S. J. Tahk, S. Y. Choi, M. H. Yoon, H. S. Lim, S. J. Kang, H. M. Yang, J. S. Park, M. Zheng, G. S. Hwang and J. H. Shin *Am J Cardiol* (2008) 102; 988-93
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18929698

Potential impact of noncontrast computed tomography as gatekeeper for myocardial perfusion positron emission tomography in patients admitted to the chest pain unit F. P. Esteves, R. Sanyal, C. A. Santana, L. Shaw and P. Raggi *Am J Cardiol* (2008) 101; 149-52
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18178397

Volumetric evaluation of coronary plaque in patients presenting with acute myocardial infarction or stable angina pectoris-a multislice computerized tomography study S. Hammer-Hansen, K. F. Kofoed, H. Kelbaek, T. Kristensen, J. T. Kuhl, J. J. Thune and L. Kober *Am Heart J* (2009) 157; 481-7
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19249418

Usefulness of 64-detector computed tomographic angiography for diagnosing in-stent restenosis in native coronary arteries H. S. Hecht, M. Zaric, V. Jelnin, L. Lubarsky, M. Prakash and G. Roubin *Am J Cardiol* (2008) 101; 820-4
<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation>

n&list_uids=18328847

Comparison of multislice computed tomography to gated single-photon emission computed tomography for imaging of healed myocardial infarcts M. M. Henneman, J. D. Schuijf, P. Dibbets-Schneider, M. P. Stokkel, R. J. van der Geest, E. E. van der Wall and J. J. Bax *Am J Cardiol* (2008) 101; 144-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18178396

Relation of a coronary artery calcium score higher than 400 to coronary stenoses detected using multidetector computed tomography and to traditional cardiovascular risk factors J. S. Ho, S. J. Fitzgerald, L. L. Stolfus, W. A. Wade, D. B. Reinhardt, C. E. Barlow and J. J. Cannaday *Am J Cardiol* (2008) 101; 1444-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18471456

Bifurcation stenting: current strategies and new devices A. Latib, A. Colombo and G. M. Sangiorgi *Heart* (2009) 95; 495-504

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18812408

Usefulness of 64-detector row computed tomography for evaluation of intracoronary stents in symptomatic patients with suspected in-stent restenosis N. Manghat, R. Van Lingen, P. Hewson, F. Syed, N. Kakani, I. Cox, C. Roobottom and G. Morgan-Hughes *Am J Cardiol* (2008) 101; 1567-73

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18489934

Association of coronary atherosclerosis detected by multislice computed tomography and traditional risk-factor assessment D. Nair, T. P. Carrigan, R. J. Curtin, Z. B. Popovic, S. Kuzmiak, P. Schoenhagen, S. D. Flamm and M. Y. Desai *Am J Cardiol* (2008) 102; 316-20

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18638593

Efficacy of culprit plaque assessment by 64-slice multidetector computed tomography to predict transient no-reflow phenomenon during percutaneous coronary intervention G. Nakazawa, K. Tanabe, Y. Onuma, S. Yachi, J. Aoki, H. Yamamoto, Y. Higashikuni, A. Yagishita, H. Nakajima and K. Hara *Am Heart J* (2008) 155; 1150-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18513532

Assessment of patients after coronary artery bypass grafting using 64-slice computed tomography I. Nazeri, P. Shahabi, M. Tehrai, B. Sharif-Kashani and A. Nazeri *Am J Cardiol* (2009) 103; 667-73

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19231331

Comparison of intravascular ultrasound to contrast-enhanced 64-slice computed tomography to assess the significance of angiographically ambiguous coronary narrowings T. Okabe, W. G. Weigold, G. S. Mintz, R. Roswell, S. Joshi, S. Y. Lee, B. Lee, D. H. Steinberg, P. Roy, T. L. Slottow, K. Smith, R. Torguson, Z. Xue, L. F. Satler, K. M. Kent, A. D. Pichard, N. J. Weissman, J. Lindsay and R. Waksman *Am J Cardiol* (2008) 102; 994-1001

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18929699

Noninvasive assessment of coronary in-stent restenosis by dual-source computed tomography T. Pflederer, M. Marwan, A. Renz, S. Bachmann, D. Ropers, A. Kuettner, K. Anders, F. Bamberg, W. G. Daniel and S. Achenbach *Am J Cardiol* (2009) 103; 812-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19268737

Clinical significance of epicardial fat measured using cardiac multislice computed tomography S. Sarin, C. Wenger, A. Marwaha, A. Qureshi, B. D. Go, C. A. Woomert, K. Clark, L. A. Nassef and J. Shirani *Am J Cardiol* (2008) 102; 767-71

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18774004

Assessment of changes in non-calcified atherosclerotic plaque volume in the left main and left anterior descending coronary arteries over time by 64-slice computed tomography M. Schmid, S. Achenbach, D. Ropers, S. Komatsu, U. Ropers, W. G. Daniel and T. Pflederer *Am J Cardiol* (2008) 101; 579-84

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18308002

Defining optimal research study design for cardiovascular imaging using computed tomography angiography as a model B. R. Shah, M. R. Patel, E. D. Peterson and P. S. Douglas *Am J Cardiol* (2008) 102; 943-8

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18805127

Computed tomographic coronary angiographic assessment of high-risk coronary anatomy in patients with suspected coronary artery disease and intermediate pretest probability T. Sheth, S. Amlani, M. L. Ellins, S. Mehta, J. Velianou, G. Cappelli, S. Yang and M. Natarajan *Am Heart J* (2008) 155; 918-23

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18440342

Impact of bivalirudin or unfractionated heparin on platelet aggregation in patients pretreated with 600 mg clopidogrel undergoing elective percutaneous coronary intervention D. Sibbing, G. Busch, S. Braun, S. Jawansky, A. Schomig, A. Kastrati, I. Ott and N. von Beckerath *Eur Heart J* (2008) 29; 1504-9

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18467322

Primary percutaneous balloon pericardiectomy for malignant pericardial effusion N. Swanson, I. Mirza, N. Wijesinghe and G. Devlin *Catheter Cardiovasc Interv* (2008) 71; 504-7

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18307242

Anatomic correlates of a normal perfusion scan using 64-slice computed tomographic coronary angiography J. M. van Werkhoven, J. D. Schuijf, J. W. Jukema, L. J. Kroft, M. P. Stokkel, P. Dibbets-Schneider, G. Pundziute, A. J. Scholte, E. E. van der Wall and J. J. Bax *Am J Cardiol* (2008) 101; 40-5

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18157963

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18157963

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18157963