

1. *Acute effects of smoking light cigarettes on coronary microvascular functions*  
Ciftci, O., et al.  
Clin Cardiol, 2009. **32**(4): p. 210-4.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19353698](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19353698)
2. *Additive prognostic value of coronary flow reserve in patients with chest pain syndrome and normal or near-normal coronary arteries*  
Sicari, R., et al.  
Am J Cardiol, 2009. **103**(5): p. 626-31.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19231324](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19231324)
3. *Association of serum adiponectin levels and coronary flow reserve in women with normal coronary angiography*  
Eroglu, S., et al.  
Eur J Cardiovasc Prev Rehabil, 2009. **16**(3): p. 290-6.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19404197](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19404197)
4. *Baseline fractional flow reserve and stent diameter predict optimal post-stent fractional flow reserve and major adverse cardiac events after bare-metal stent deployment*  
Samady, H., et al.  
JACC Cardiovasc Interv, 2009. **2**(4): p. 357-63.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19463450](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19463450)
5. *Clinical outcome in patients with intermediate stenosis of left anterior descending coronary artery after deferral of revascularization on the basis of noninvasive coronary flow reserve measurement*  
D'Andrea, A., et al.  
Echocardiography, 2009. **26**(4): p. 431-40.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19054024](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19054024)
6. *Comparison of non-invasive multi-slice computed tomography coronary angiography versus invasive coronary angiography and fractional flow reserve for the evaluation of men with known coronary artery disease*  
van Werkhoven, J.M., et al.  
Am J Cardiol, 2009. **104**(5): p. 653-6.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19699340](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19699340)

7. *Comparison of the intracoronary continuous infusion method using a microcatheter and the intravenous continuous adenosine infusion method for inducing maximal hyperemia for fractional flow reserve measurement*

Yoon, M.H., et al.

Am Heart J, 2009. **157**(6): p. 1050-6.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19464416](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19464416)

8. *Coronary flow reserve is impaired in hypertensive patients with subclinical renal damage*

Bezante, G.P., et al.

Am J Hypertens, 2009. **22**(2): p. 191-6.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19151691](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19151691)

9. *Coronary flow velocity reserve and aortic distensibility indices in hypertensive patients with hypercholesterolaemia and normal epicardial coronary arteries*

Nemes, A., T. Forster, and M. Csanady

Clin Exp Hypertens, 2009. **31**(4): p. 380-7.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19811366](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19811366)

10. *Coronary microvascular resistance index immediately after primary percutaneous coronary intervention as a predictor of the transmural extent of infarction in patients with ST-segment elevation anterior acute myocardial infarction*

Kitabata, H., et al.

JACC Cardiovasc Imaging, 2009. **2**(3): p. 263-72.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19356570](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19356570)

11. *The delta fractional flow reserve can predict lesion severity and long-term prognosis*

Kocaman, S.A., et al.

Atherosclerosis, 2009. **203**(1): p. 178-84.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18644595](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18644595)

12. *Endothelial and non-endothelial coronary blood flow reserve and left ventricular dysfunction in systemic hypertension*  
Rocha, A.M., et al.  
Clinics (Sao Paulo), 2009. **64**(4): p. 327-35.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19488591](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19488591)
13. *Endothelial-mediated coronary flow reserve in patients with mild thyroid hormone deficiency*  
Biondi, B., et al.  
Eur J Endocrinol, 2009. **161**(2): p. 323-9.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19423563](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19423563)
14. *Evaluation of coronary microvascular function in patients with end-stage renal disease, and renal allograft recipients*  
Bozbas, H., et al.  
Atherosclerosis, 2009. **202**(2): p. 498-504.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18550064](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18550064)
15. *Evaluation of intermediate coronary stenosis with intravascular ultrasound and fractional flow reserve: Its use and abuse*  
Magni, V., A. Chieffo, and A. Colombo  
Catheter Cardiovasc Interv, 2009. **73**(4): p. 441-8.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19133668](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19133668)
16. *Features of cardiac allograft coronary endothelial dysfunction*  
Raichlin, E., et al.  
Am J Cardiol, 2009. **103**(8): p. 1154-8.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19361606](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19361606)
17. *Fractional flow reserve versus angiography for guiding percutaneous coronary intervention*  
Tonino, P.A., et al.  
N Engl J Med, 2009. **360**(3): p. 213-24.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19144937](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19144937)

18. *The impact of multiple vessel disease on fractional flow reserve*  
Sahinarslan, A., et al.  
Acta Cardiol, 2009. **64**(1): p. 79-83.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19317302](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19317302)
19. *The impact of sleep deprivation on the coronary circulation*  
Sekine, T., et al.  
Int J Cardiol, 2009.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19203808](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19203808)
20. *Impairment of coronary flow reserve in aortic stenosis*  
Garcia, D., et al.  
J Appl Physiol, 2009. **106**(1): p. 113-21.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18974370](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18974370)
21. *Improvement of fractional flow reserve and collateral flow by treatment with external counterpulsation (Art.Net.-2 Trial)*  
Buschmann, E.E., et al.  
Eur J Clin Invest, 2009. **39**(10): p. 866-75.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19572918](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19572918)
22. *Influence of abnormal glucose metabolism on coronary microvascular function after a recent myocardial infarction*  
Logstrup, B.B., et al.  
JACC Cardiovasc Imaging, 2009. **2**(10): p. 1159-66.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19833304](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19833304)
23. *Inhomogeneous vasomotor effects of moderate selective and non-selective endothelin-receptor blockade in stable coronary artery disease*  
Wexberg, P., et al.  
Heart, 2009. **95**(15): p. 1258-64.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19414437](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19414437)

24. *Long-term clinical outcome after fractional flow reserve-guided treatment in patients with angiographically equivocal left main coronary artery stenosis*  
Hamilos, M., et al.  
Circulation, 2009. **120**(15): p. 1505-12.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19786633](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19786633)
25. *Long-term prognostic role of coronary flow velocity reserve in patients with aortic valve stenosis - insights from the SZEGED Study*  
Nemes, A., et al.  
Clin Physiol Funct Imaging, 2009. **29**(6): p. 447-52.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19712079](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19712079)
26. *Myocardial bridging in absence of coronary artery disease: proposal of a new classification based on clinical-angiographic data and long-term follow-up*  
Schwarz, E.R., et al.  
Cardiology, 2009. **112**(1): p. 13-21.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18577881](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18577881)
27. *Non-invasive coronary flow reserve after successful primary angioplasty for acute anterior myocardial infarction is an independent predictor of left ventricular recovery and in-hospital cardiac events*  
Meimoun, P., et al.  
J Am Soc Echocardiogr, 2009. **22**(9): p. 1071-9.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19647405](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19647405)
28. *Patients with coronary slow flow phenomenon demonstrate normal myocardial blood flow and arterial wave reflection between acute episodes*  
Sharman, J.E., et al.  
Int J Cardiol, 2009. **131**(3): p. 321-5.  
[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18582970](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18582970)
29. *Prognostic correlates of combined coronary flow reserve assessment on left anterior descending and right coronary artery in patients with negative stress echocardiography by wall motion criteria*  
Cortigiani, L., et al.  
Heart, 2009. **95**(17): p. 1423-8.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19414438](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19414438)

30. *Prognostic value of transthoracic coronary flow reserve in medically treated patients with proximal left anterior descending artery stenosis of intermediate severity*

Meimoun, P., et al.

Eur J Echocardiogr, 2009. **10**(1): p. 127-32.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=18579502](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18579502)

31. *Relationship between functional exercise capacity and functional stenosis in patients with stable angina and intermediate coronary stenosis*

Tanaka, S., et al.

Circ J, 2009. **73**(12): p. 2308-14.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19838006](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19838006)

32. *Relationship between hyperglycemia and coronary vascular resistance in non-diabetic patients*

Ichiki, H., et al.

Int J Cardiol, 2009.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19147243](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19147243)

33. *The reliability of fractional flow reserve measurement in patients with diabetes mellitus*

Sahinarslan, A., et al.

Coron Artery Dis, 2009. **20**(5): p. 317-21.

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19444091](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19444091)